

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
14 October 2004 (14.10.2004)

PCT

(10) International Publication Number  
WO 2004/087704 A1

(51) International Patent Classification<sup>7</sup>: C07D 471/04,  
A61K 31/437, A61P 25/04

(74) Agent: GLOBAL INTELLECTUAL PROPERTY; As-  
traZeneca AB, S-151 85 Södertälje (SE).

(21) International Application Number:  
PCT/SE2004/000472

(22) International Filing Date: 26 March 2004 (26.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0300908-1 31 March 2003 (31.03.2003) SE

(71) Applicant (for all designated States except US): AS-  
TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): WEL, Zhongy-  
ong [CA/CA]; AstraZeneca R & D Montreal, 7171  
Frederick-Banting, St Laurent, Québec H4S 1Z9 (CA).  
DOLAINE, Regis [FR/CA]; AstraZeneca R & D Mon-  
treal, 7171 Frederick-Banting, St Laurent, Québec H4S  
1Z9 (CA). WALPOLE, Christopher [GB/CA]; As-  
traZeneca R & D Montreal, 7171 Frederick-Banting, St  
Laurent, Québec H4S 1Z9 (CA). YANG, Hua [CA/CA];  
AstraZeneca R & D Montreal, 7171 Frederick-Banting, St  
Laurent, Québec H4S 1Z9 (CA).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,  
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,  
ZW.


(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-  
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,  
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: AZAINDOLE DERIVATIVES, PREPARATIONS THEREOF, USES THEREOF AND COMPOSITIONS CONTAIN-  
ING THEM



(57) Abstract: Compounds of formula I or pharmaceutically acceptable salts thereof Formula (I) wherein Ar, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>a</sup>, R<sup>b</sup> and X are as defined in the specifications as well as salts and pharmaceutical compositions including the compounds are prepared. They are useful in therapy, in particular in the management of pain.

WO 2004/087704 A1

JC20 Rec'd PCT/PTO 2 6 SEP 2005

**AZAINDOLE DERIVATIVES, PREPARATIONS THEREOF, USES THEREOF  
AND COMPOSITIONS CONTAINING THEM**

**5    BACKGROUND OF THE INVENTION**

**1. Field of the invention**

The invention is related to compounds which are CB<sub>1</sub>/CB<sub>2</sub> receptor ligands, pharmaceutical compositions containing these compounds, manufacturing processes thereof and uses thereof, and more particularly to compounds that are CB<sub>1</sub>/CB<sub>2</sub> receptor agonists.

**10    2. Discussion of Relevant Technology**

Pain management has been an important field of study for many years. It has been well known that cannabinoid receptor (e.g., CB<sub>1</sub> receptors, CB<sub>2</sub> receptors) ligands, especially agonists produce relief of pain in a variety of animal models by interacting with CB<sub>1</sub> and/or CB<sub>2</sub> receptors. Generally, CB<sub>1</sub> receptors are located predominately in the central nervous  
**15** system, whereas CB<sub>2</sub> receptors are located primarily in the periphery and are primarily restricted to the cells and tissues derived from the immune system.

While the conventional CB<sub>1</sub> receptor agonists and CB<sub>1</sub>/CB<sub>2</sub> receptor agonists, such as tetrahydrocannabinol (THC) and *Cannabis*-related drugs, are highly effective in anti-nociception models in animals, they tend to exert many undesired CNS (central nerve system)  
**20** side-effects, e.g., psychoactive side effects and the abuse potential of *Cannabis*-related drugs.

Therefore, there is a need for new CB<sub>1</sub>/CB<sub>2</sub> receptor ligands such as agonists useful in managing pain or treating other related symptoms or diseases with reduced or minimal undesirable CNS side-effects.

**DISCLOSURE OF THE INVENTION**

**25**        The present invention provides CB<sub>1</sub>/CB<sub>2</sub> receptor ligands which are useful in treating pain and other related symptoms or diseases.

**Definitions**

Unless specified otherwise within this specification, the nomenclature used in this specification generally follows the examples and rules stated in *Nomenclature of Organic*  
**30** *Chemistry, Sections A, B, C, D, E, F, and H*, Pergamon Press, Oxford, 1979, which is incorporated by references herein for its exemplary chemical structure names and rules on naming chemical structures. Optionally, a name of a compound may be generated using a chemical naming program: ACD/ChemSketch, Version 5.09/September 2001, Advanced Chemistry Development, Inc., Toronto, Canada.

- 2 -

"CB<sub>1</sub>/CB<sub>2</sub> receptors" means CB<sub>1</sub> and/or CB<sub>2</sub> receptors.

The term "C<sub>m-n</sub>" or "C<sub>m-n</sub> group" used alone or as a prefix, refers to any group having m to n carbon atoms, and having 0 to n multivalent heteroatoms selected from O, S, N and P, wherein m and n are 0 or positive integers, and n>m. For example, "C<sub>1-6</sub>" would refer to a chemical group having 1 to 6 carbon atoms, and having 0 to 6 multivalent heteroatoms selected from O, S, N and P.

The term "hydrocarbon" used alone or as a suffix or prefix, refers to any structure comprising only carbon and hydrogen atoms up to 14 carbon atoms.

The term "hydrocarbon radical" or "hydrocarbonyl" used alone or as a suffix or prefix, refers to any structure as a result of removing one or more hydrogens from a hydrocarbon.

The term "alkyl" used alone or as a suffix or prefix, refers to monovalent straight or branched chain hydrocarbon radicals comprising 1 to about 12 carbon atoms. Unless otherwise specified, "alkyl" general includes both saturated alkyl and unsaturated alkyl.

The term "alkylene" used alone or as suffix or prefix, refers to divalent straight or branched chain hydrocarbon radicals comprising 1 to about 12 carbon atoms, which serves to links two structures together.

The term "alkenyl" used alone or as suffix or prefix, refers to a monovalent straight or branched chain hydrocarbon radical having at least one carbon-carbon double bond and comprising at least 2 up to about 12 carbon atoms.

The term "alkynyl" used alone or as suffix or prefix, refers to a monovalent straight or branched chain hydrocarbon radical having at least one carbon-carbon triple bond and comprising at least 2 up to about 12 carbon atoms.

The term "cycloalkyl," used alone or as suffix or prefix, refers to a monovalent ring-containing hydrocarbon radical comprising at least 3 up to about 12 carbon atoms.

The term "cycloalkenyl" used alone or as suffix or prefix, refers to a monovalent ring-containing hydrocarbon radical having at least one carbon-carbon double bond and comprising at least 3 up to about 12 carbon atoms.

The term "cycloalkynyl" used alone or as suffix or prefix, refers to a monovalent ring-containing hydrocarbon radical having at least one carbon-carbon triple bond and comprising about 7 up to about 12 carbon atoms.

The term "aryl" used alone or as suffix or prefix, refers to a monovalent hydrocarbon radical having one or more polyunsaturated carbon rings having aromatic character, (e.g., 4n + 2 delocalized electrons) and comprising 5 up to about 14 carbon atoms.

The term "arylene" used alone or as suffix or prefix, refers to a divalent hydrocarbon radical having one or more polyunsaturated carbon rings having aromatic character, (*e.g.*,  $4n + 2$  delocalized electrons) and comprising 5 up to about 14 carbon atoms, which serves to links two structures together.

- 5       The term "heterocycle" used alone or as a suffix or prefix, refers to a ring-containing structure or molecule having one or more multivalent heteroatoms, independently selected from N, O, P and S, as a part of the ring structure and including at least 3 and up to about 20 atoms in the ring(s). Heterocycle may be saturated or unsaturated, containing one or more double bonds, and heterocycle may contain more than one ring. When a heterocycle contains  
10 more than one ring, the rings may be fused or unfused. Fused rings generally refer to at least two rings share two atoms therebetween. Heterocycle may have aromatic character or may not have aromatic character.

- The term "heteroalkyl" used alone or as a suffix or prefix, refers to a radical formed as a result of replacing one or more carbon atom of an alkyl with one or more heteroatoms  
15 selected from N, O, P and S.

- The term "heteroaromatic" used alone or as a suffix or prefix, refers to a ring-containing structure or molecule having one or more multivalent heteroatoms, independently selected from N, O, P and S, as a part of the ring structure and including at least 3 and up to about 20 atoms in the ring(s), wherein the ring-containing structure or molecule has an  
20 aromatic character (*e.g.*,  $4n + 2$  delocalized electrons).

      The term "heterocyclic group," "heterocyclic moiety," "heterocyclic," or "heterocyclo" used alone or as a suffix or prefix, refers to a radical derived from a heterocycle by removing one or more hydrogens therefrom.

- The term "heterocyclyl" used alone or as a suffix or prefix, refers a monovalent radical  
25 derived from a heterocycle by removing one hydrogen therefrom.

      The term "heterocyclylene" used alone or as a suffix or prefix, refers to a divalent radical derived from a heterocycle by removing two hydrogens therefrom, which serves to links two structures together.

- The term "heteroaryl" used alone or as a suffix or prefix, refers to a heterocyclyl  
30 having aromatic character.

      The term "heterocylcoalkyl" used alone or as a suffix or prefix, refers to a heterocyclyl that does not have aromatic character.

The term "heteroarylene" used alone or as a suffix or prefix, refers to a heterocyclylene having aromatic character.

The term "heterocycloalkylene" used alone or as a suffix or prefix, refers to a heterocyclylene that does not have aromatic character.

5 The term "six-membered" used as prefix refers to a group having a ring that contains six ring atoms.

The term "five-membered" used as prefix refers to a group having a ring that contains five ring atoms.

A five-membered ring heteroaryl is a heteroaryl with a ring having five ring atoms  
10 wherein 1, 2 or 3 ring atoms are independently selected from N, O and S.

Exemplary five-membered ring heteroaryls are thienyl, furyl, pyrrolyl, imidazolyl, thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-triazolyl, 1,2,4-thiadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-triazolyl, 1,3,4-thiadiazolyl, and 1,3,4-oxadiazolyl.

15 A six-membered ring heteroaryl is a heteroaryl with a ring having six ring atoms wherein 1, 2 or 3 ring atoms are independently selected from N, O and S.

Exemplary six-membered ring heteroaryls are pyridyl, pyrazinyl, pyrimidinyl, triazinyl and pyridazinyl.

The term "substituted" used as a prefix refers to a structure, molecule or group,  
20 wherein one or more hydrogens are replaced with one or more C<sub>1-12</sub>hydrocarbon groups, or one or more chemical groups containing one or more heteroatoms selected from N, O, S, F, Cl, Br, I, and P. Exemplary chemical groups containing one or more heteroatoms include heterocyclyl, -NO<sub>2</sub>, -OR, -Cl, -Br, -I, -F, -CF<sub>3</sub>, -C(=O)R, -C(=O)OH, -NH<sub>2</sub>, -SH, -NHR, -NR<sub>2</sub>, -SR, -SO<sub>3</sub>H, -SO<sub>2</sub>R, -S(=O)R, -CN, -OH, -C(=O)OR, -C(=O)NR<sub>2</sub>, -NRC(=O)R, oxo  
25 (=O), imino (=NR), thio (=S), and oximino (=N-OR), wherein each "R" is a C<sub>1-12</sub>hydrocarbyl. For example, substituted phenyl may refer to nitrophenyl, pyridylphenyl, methoxyphenyl, chlorophenyl, aminophenyl, etc., wherein the nitro, pyridyl, methoxy, chloro, and amino groups may replace any suitable hydrogen on the phenyl ring.

The term "substituted" used as a suffix of a first structure, molecule or group,  
30 followed by one or more names of chemical groups refers to a second structure, molecule or group, which is a result of replacing one or more hydrogens of the first structure, molecule or group with the one or more named chemical groups. For example, a "phenyl substituted by nitro" refers to nitrophenyl.

- 5 -

The term "optionally substituted" refers to both groups, structures, or molecules that are substituted and those that are not substituted.

Heterocycle includes, for example, monocyclic heterocycles such as: aziridine, oxirane, thiirane, azetidine, oxetane, thietane, pyrrolidine, pyrroline, imidazolidine, pyrazolidine, pyrazoline, dioxolane, sulfolane, 2,3-dihydrofuran, 2,5-dihydrofuran, tetrahydrofuran, thiophane, piperidine, 1,2,3,6-tetrahydro-pyridine, piperazine, morpholine, thiomorpholine, pyran, thiopyran, 2,3-dihydropyran, tetrahydropyran, 1,4-dihydropyridine, 1,4-dioxane, 1,3-dioxane, dioxane, homopiperidine, 2,3,4,7-tetrahydro-1*H*-azepine, homopiperazine, 1,3-dioxepane, 4,7-dihydro-1,3-dioxepin, and hexamethylene oxide.

In addition, heterocycle includes aromatic heterocycles, for example, pyridine, pyrazine, pyrimidine, pyridazine, thiophene, furan, furazan, pyrrole, imidazole, thiazole, oxazole, pyrazole, isothiazole, isoxazole, 1,2,3-triazole, tetrazole, 1,2,3-thiadiazole, 1,2,3-oxadiazole, 1,2,4-triazole, 1,2,4-thiadiazole, 1,2,4-oxadiazole, 1,3,4-triazole, 1,3,4-thiadiazole, and 1,3,4-oxadiazole.

Additionally, heterocycle encompass polycyclic heterocycles, for example, indole, indoline, isoindoline, quinoline, tetrahydroquinoline, isoquinoline, tetrahydroisoquinoline, 1,4-benzodioxan, coumarin, dihydrocoumarin, benzofuran, 2,3-dihydrobenzofuran, isobenzofuran, chromene, chroman, isochroman, xanthene, phenoxathiin, thianthrene, indolizine, isoindole, indazole, purine, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, phenanthridine, perimidine, phenanthroline, phenazine, phenothiazine, phenoxazine, 1,2-benzisoxazole, benzothiophene, benzoxazole, benzthiazole, benzimidazole, benztriazole, thioxanthine, carbazole, carboline, acridine, pyrolizidine, and quinolizidine.

In addition to the polycyclic heterocycles described above, heterocycle includes polycyclic heterocycles wherein the ring fusion between two or more rings includes more than one bond common to both rings and more than two atoms common to both rings. Examples of such bridged heterocycles include quinuclidine, diazabicyclo[2.2.1]heptane and 7-oxabicyclo[2.2.1]heptane.

Heterocyclyl includes, for example, monocyclic heterocyclyls, such as: aziridinyl, oxiranyl, thiiranyl, azetidiny, oxetanyl, thietanyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, pyrazolidinyl, pyrazolinyl, dioxolanyl, sulfolanyl, 2,3-dihydrofuranyl, 2,5-dihydrofuranyl, tetrahydrofuranyl, thiophanyl, piperidinyl, 1,2,3,6-tetrahydro-pyridinyl, piperazinyl, morpholinyl, thiomorpholinyl, pyranyl, thiopyranyl, 2,3-dihydropyranyl, tetrahydropyranyl, 1,4-dihydropyridinyl, 1,4-dioxanyl, 1,3-dioxanyl, dioxanyl, homopiperidinyl, 2,3,4,7-

tetrahydro-1*H*-azepinyl, homopiperazinyl, 1,3-dioxepanyl, 4,7-dihydro-1,3-dioxepinyl, and hexamethylene oxidyl.

In addition, heterocyclyl includes aromatic heterocyclyls or heteroaryl, for example, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, thienyl, furyl, furazanyl, pyrrolyl, imidazolyl, 5 thiazolyl, oxazolyl, pyrazolyl, isothiazolyl, isoxazolyl, 1,2,3-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-triazolyl, 1,2,4-thiadiazolyl, 1,2,4-oxadiazolyl, 1,3,4-triazolyl, 1,3,4-thiadiazolyl, and 1,3,4 oxadiazolyl.

Additionally, heterocyclyl encompasses polycyclic heterocyclyls (including both aromatic or non-aromatic), for example, indolyl, indolinyl, isoindolinyl, quinolinyl, 10 tetrahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, 1,4-benzodioxanyl, coumarinyl, dihydrocoumarinyl, benzofuranyl, 2,3-dihydrobenzofuranyl, isobenzofuranyl, chromenyl, chromanyl, isochromanyl, xanthenyl, phenoxathiinyl, thianthrenyl, indolizinyl, isoindolyl, indazolyl, purinyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, phenanthridinyl, perimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, 15 phenoxazinyl, 1,2-benzisoxazolyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benzimidazolyl, benztriazolyl, thioxanthinyl, carbazolyl, carbolinyl, acridinyl, pyrolizidinyl, and quinolizidinyl.

In addition to the polycyclic heterocyclyls described above, heterocyclyl includes polycyclic heterocyclyls wherein the ring fusion between two or more rings includes more 20 than one bond common to both rings and more than two atoms common to both rings. Examples of such bridged heterocycles include quinuclidinyl, diazabicyclo[2.2.1]heptyl; and 7-oxabicyclo[2.2.1]heptyl.

The term "alkoxy" used alone or as a suffix or prefix, refers to radicals of the general formula -O-R, wherein -R is selected from a hydrocarbon radical. Exemplary alkoxy 25 includes methoxy, ethoxy, propoxy, isopropoxy, butoxy, t-butoxy, isobutoxy, cyclopropylmethoxy, allyloxy, and propargyloxy.

The term "aryloxy" used alone or as suffix or prefix, refers to radicals of the general formula -O-Ar, wherein -Ar is an aryl.

The term "heteroaryloxy" used alone or as suffix or prefix, refers to radicals of the 30 general formula -O-Ar', wherein -Ar' is a heteroaryl.

The term "amine" or "amino" used alone or as a suffix or prefix, refers to radicals of the general formula -NRR', wherein R and R' are independently selected from hydrogen or a hydrocarbon radical.

- 7 -

"Acyl" used alone, as a prefix or suffix, means  $-C(=O)-R$ , wherein  $-R$  is an optionally substituted hydrocarbyl, hydrogen, amino or alkoxy. Acyl groups include, for example, acetyl, propionyl, benzoyl, phenyl acetyl, carboethoxy, and dimethylcarbamoyl.

Halogen includes fluorine, chlorine, bromine and iodine.

5 "Halogenated," used as a prefix of a group, means one or more hydrogens on the group is replaced with one or more halogens.

"RT" or "rt" means room temperature.

A first ring group being "fused" with a second ring group means the first ring and the second ring share at least two atoms therebetween.

10 "Link," "linked," or "linking," unless otherwise specified, means covalently linked or bonded.


When a first group, structure, or atom is "directly connected" to a second group, structure or atom, at least one atom of the first group, structure or atom forms a chemical bond with at least one atom of the second group, structure or atom.

15 "Saturated carbon" means a carbon atom in a structure, molecule or group wherein all the bonds connected to this carbon atom are single bond. In other words, there is no double or triple bonds connected to this carbon atom and this carbon atom generally adopts an  $sp^3$  atomic orbital hybridization.

"Unsaturated carbon" means a carbon atom in a structure, molecule or group wherein at least one bond connected to this carbon atom is not a single bond. In other words, there is at least one double or triple bond connected to this carbon atom and this carbon atom generally adopts a  $sp$  or  $sp^2$  atomic orbital hybridization.

#### Description of Preferred Embodiments

25 In one aspect, the invention provides a compound of formula I, a pharmaceutically acceptable salt thereof, diastereomers, enantiomers, or mixtures thereof:



**I**

wherein

$R^1$  is a  $C_{1-12}$  group;



- 8 -

X is a C<sub>1-10</sub> divalent group that separates groups connected thereto by one or two saturated carbons;

Ar is C<sub>4-12</sub> divalent aromatic group;

R<sup>2</sup> is optionally substituted C<sub>1-6</sub>hydrocarbaryl, optionally substituted C<sub>6-10</sub>aryl, or  
5 optionally substituted C<sub>3-6</sub>heteroaryl;

R<sup>3</sup> is a C<sub>1-12</sub> group, wherein the atom of R<sup>3</sup> that is directly connected to the six-membered ring of formula I is a nitrogen, or an unsaturated carbon, wherein the unsaturated carbon is connected to an oxygen through a double bond; and

R<sup>a</sup> and R<sup>b</sup> are -R, -NO<sub>2</sub>, -OR, -Cl, -Br, -I, -F, -CF<sub>3</sub>, -C(=O)R, -C(=O)OH, -NH<sub>2</sub>, -SH, -  
10 NHR, -NR<sub>2</sub>, -SR, -SO<sub>3</sub>H, -SO<sub>2</sub>R, -S(=O)R, -CN, -OH, -C(=O)OR, or -NRC(=O)R, wherein R is independently -H or C<sub>1-6</sub> hydrocarbaryl.


Particularly, the compounds of the present invention are those of formula I, wherein

R<sup>1</sup> is optionally substituted C<sub>1-10</sub> hydrocarbaryl; optionally substituted C<sub>1-10</sub>acyl;  
optionally substituted C<sub>4-8</sub>heteroaryl-C(=O)-; R<sup>4</sup>R<sup>5</sup>N-C<sub>1-6</sub>alkyl; R<sup>4</sup>R<sup>5</sup>NC(=O)-C<sub>1-6</sub>alkyl; R<sup>4</sup>O-  
15 C<sub>1-6</sub> alkyl; R<sup>4</sup>OC(=O)-C<sub>1-6</sub>alkyl; R<sup>4</sup>C(=O)-C<sub>1-6</sub>alkyl; R<sup>4</sup>C(=O)NR<sup>4</sup>-C<sub>1-6</sub>alkyl; R<sup>4</sup>R<sup>5</sup>NSO<sub>2</sub>-C<sub>1-6</sub>alkyl; R<sup>4</sup>CSO<sub>2</sub>N(R<sup>5</sup>)-C<sub>1-6</sub>alkyl; R<sup>4</sup>R<sup>5</sup>NC(=O)N(R<sup>6</sup>)-C<sub>1-6</sub>alkyl; R<sup>4</sup>R<sup>5</sup>NSO<sub>2</sub>N(R<sup>6</sup>)-C<sub>1-6</sub>alkyl;  
optionally substituted aryl-C<sub>1-6</sub>alkyl; optionally substituted aryl-C(=O)-C<sub>1-6</sub>alkyl; optionally substituted heterocyclyl-C<sub>1-6</sub>alkyl; optionally substituted heterocyclyl-C(=O)-C<sub>1-6</sub>alkyl; and C<sub>1-10</sub>hydrocarbaryl-amino;

20 wherein R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> are independently selected from -H, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, or a divalent C<sub>1-6</sub>group that together with another divalent C<sub>1-6</sub>group forms a portion of a ring;

R<sup>2</sup> is optionally substituted C<sub>1-6</sub>hydrocarbaryl, optionally substituted C<sub>6-10</sub>aryl, or optionally substituted C<sub>3-6</sub>heteroaryl;

25 R<sup>3</sup> is selected from:



wherein

- 9 -

$R^7$  is selected from -H, optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl;

$R^{10}$ ,  $R^{11}$ ,  $R^{12}$  and  $R^{13}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl; and  $R^a$  and  $R^b$  are hydrogen.


More particularly, the compounds of the present invention are those of formula I, wherein  $R^1$  is selected from  $C_{1-8}$ alkyl;  $C_{2-8}$ alkenyl;  $C_{2-8}$ alkynyl; optionally substituted aryl- $C_{1-6}$ alkyl;  $R^4R^5NC_{1-6}$ alkyl;  $R^4OC_{1-6}$ alkyl;  $C_{3-6}$ cycloalkyl- $C_{1-6}$ alkyl; optionally substituted  $C_{3-6}$ heterocycloalkyl- $C_{1-6}$ alkyl;  $C_{1-6}$ alkyl- $C_{6-8}$ aryl;  $C_{1-6}$ alkyl-C(=O)-;  $C_{6-8}$ aryl-C(=O)-;  $C_{3-6}$ heteroaryl-C(=O)-; or optionally substituted  $C_{3-6}$ heteroaryl- $C_{1-6}$ alkyl;

wherein  $R^2$  is selected from  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyl substituted by at least one fluorine,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkenyl substituted by at least one fluorine,  $C_{2-6}$ alkynyl,  $C_{2-6}$ alkynyl substituted by at least one fluorine, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$ aryl, and optionally substituted  $C_{3-6}$ heteroaryl;

$R^4$ ,  $R^5$  and  $R^6$  are independently selected from the group consisting of -H,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl, and a divalent  $C_{1-6}$ group that together with another divalent  $C_{1-6}$ group forms a portion of a ring; and

X is selected from the group consisting of  $-NR^6-$ ,  $-CH_2-CH_2-$ ,  $-CH=CH-$ ,  $-O-$ ,  $-C(R^8)(R^9)-$ , and  $-S(O)_q-$ , wherein q is 0, 1 or 2, wherein  $R^8$  and  $R^9$  are independently  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{1-6}$ alkoxy, -OH, or -H; at most one of  $R^8$  and  $R^9$  is -OH.

$R^3$  is selected from:



25 wherein

- 10 -

$R^7$  is selected from -H, optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl;

$R^{10}$ ,  $R^{11}$ ,  $R^{12}$  and  $R^{13}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl; and

$R^a$  and  $R^b$  are hydrogen.

In a more particular embodiment, the compounds of the present invention are those of formula I, wherein


$R^1$  is selected from  $C_{1-6}$ alkyl;  $C_{2-6}$ alkenyl;  $C_{2-6}$ alkynyl; optionally substituted  $C_{3-6}$ cycloalkylmethyl; optionally substituted  $C_{3-6}$ heterocycloalkylmethyl;

X is  $-CH_2-$ ;

Ar is phenylene or pyridylene;

$R^2$  is selected from  $-CH_3$ ,  $-CH_2CH_3$ ,  $-CH(CH_3)_2$ ,  $-CH_2CF_3$ ,  $CF_3$ , cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyridyl and phenyl; and

$R^3$  is selected from:



wherein,  $R^7$  is selected from -H and methyl;  $R^{10}$  and  $R^{11}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl.

In another more particular embodiment, the compounds of the present invention are those of formula I, wherein

$R^1$  is selected from  $C_{1-6}$ alkyl;  $C_{2-6}$ alkenyl;  $C_{2-6}$ alkynyl; optionally substituted  $C_{3-6}$ cycloalkylmethyl; optionally substituted  $C_{3-6}$ heterocycloalkylmethyl;


X is  $-CH_2-$ ;

Ar is selected from the group consisting of an optionally substituted *para*-arylene; an optionally substituted a six-membered *para*-heteroarylene;

$R^2$  is selected from  $-CH_3$ ,  $-CH_2CH_3$ ,  $-CH(CH_3)_2$ ,  $-CH_2CF_3$ ,  $CF_3$ , cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyridyl and phenyl; and

$R^3$  is selected from:

- 11 -



wherein,  $R^7$  is selected from -H and methyl;  $R^{10}$  and  $R^{11}$  are selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl.

Most particularly, the compounds of the present invention are those of formula I, wherein


$R^1$  is selected from optionally substituted  $C_{3-6}$ cycloalkylmethyl; and optionally substituted  $C_{3-6}$ heterocycloalkylmethyl;

10  $X$  is  $-CH_2-$ ;

$Ar$  is *para*-phenylene or *para*-pyridylene;

$R^2$  is methyl, or ethyl; and

$R^3$  is selected from:



15 wherein,  $R^7$  is selected from -H and methyl;  $R^{10}$  and  $R^{11}$  are selected from  $C_{1-6}$ alkyl,  $C_{3-6}$ cycloalkyl, phenyl optionally substituted with halogen, nitro,  $C_{1-3}$ alkyl,  $-COOR^{14}$ ,  $-OH$ , cyano, trifluoromethyl,  $C_{1-3}$ alkyloxy;  $C_{3-6}$ heteroaryl optionally substituted with halogen, nitro,  $C_{1-3}$ alkyl,  $-COOR^{14}$ ,  $-OH$ , cyano, trifluoromethyl,  $C_{1-3}$ alkyloxy, wherein  $R^{14}$  is a  $C_{1-3}$ alkyl.

It will be understood that when compounds of the present invention contain one or  
 20 more chiral centers, the compounds of the invention may exist in, and be isolated as, enantiomeric or diastereomeric forms, or as a racemic mixture. The present invention includes any possible enantiomers, diastereomers, racemates or mixtures thereof, of a compound of Formula I. The optically active forms of the compound of the invention may be prepared, for example, by chiral chromatographic separation of a racemate, by synthesis from  
 25 optically active starting materials or by asymmetric synthesis based on the procedures described thereafter.

It will also be appreciated that certain compounds of the present invention may exist as geometrical isomers, for example E and Z isomers of alkenes. The present invention

- 12 -

includes any geometrical isomer of a compound of Formula I. It will further be understood that the present invention encompasses tautomers of the compounds of the formula I.

It will also be understood that certain compounds of the present invention may exist in solvated, for example hydrated, as well as unsolvated forms. It will further be understood that  
5 the present invention encompasses all such solvated forms of the compounds of the formula I.

Within the scope of the invention are also salts of the compounds of the formula I. Generally, pharmaceutically acceptable salts of compounds of the present invention may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound, for example an alkyl amine with a suitable acid, for example,  
10 HCl or acetic acid, to afford a physiologically acceptable anion. It may also be possible to make a corresponding alkali metal (such as sodium, potassium, or lithium) or an alkaline earth metal (such as a calcium) salt by treating a compound of the present invention having a suitably acidic proton, such as a carboxylic acid or a phenol with one equivalent of an alkali metal or alkaline earth metal hydroxide or alkoxide (such as the ethoxide or methoxide), or a  
15 suitably basic organic amine (such as choline or meglumine) in an aqueous medium, followed by conventional purification techniques.

In one embodiment, the compound of formula I above may be converted to a pharmaceutically acceptable salt or solvate thereof, particularly, an acid addition salt such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate,  
20 methanesulphonate or *p*-toluenesulphonate.

We have discovered that the compounds of the invention have activity as pharmaceuticals, in particular as modulators or ligands such as agonists, partial agonists, inverse agonist or antagonists of CB<sub>1</sub>/CB<sub>2</sub> receptors. More particularly, the compounds of the invention exhibit selective activity as agonists of the CB<sub>1</sub>/CB<sub>2</sub> receptors, and are useful in the  
25 relief of pain, particularly chronic pain, e.g., chronic inflammatory pain, neuropathic pain, back pain, cancer pain and visceral pain. Compounds of the present invention will also be useful in treating acute pain, anxiety disorders, gastrointestinal disorders, cardiovascular disorders, multiple sclerosis, Parkinson's disease, Huntington's chorea, Alzheimer's disease and/or cancers of the immune system or cells thereof. Additionally, compounds of the present  
30 invention are useful in other disease states in which degeneration or dysfunction of CB<sub>1</sub>/CB<sub>2</sub> receptors is present or implicated.

Thus, the invention provides a compound of formula I, or pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.

- 13 -

In a further aspect, the present invention provides the use of a compound of formula I, or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The term "therapeutic" and "therapeutically" should be construed accordingly. The term "therapy" within the context of the present invention further encompasses to administer an effective amount of a compound of the present invention, to mitigate either a pre-existing disease state, acute or chronic, or a recurring condition. This definition also encompasses prophylactic therapies for prevention of recurring conditions and continued therapy for chronic disorders.

The compounds of the present invention are useful in therapy, especially for the therapy of various pain conditions including, but not limited to: acute pain, chronic pain, neuropathic pain, back pain, cancer pain, and visceral pain.

In use for therapy in a warm-blooded animal such as a human, the compound of the invention may be administered in the form of a conventional pharmaceutical composition by any route including orally, intramuscularly, subcutaneously, topically, intranasally, intraperitoneally, intrathoracically, intravenously, epidurally, intrathecally, intracerebroventricularly and by injection into the joints.

In one embodiment of the invention, the route of administration may be orally, intravenously or intramuscularly.

The dosage will depend on the route of administration, the severity of the disease, age and weight of the patient and other factors normally considered by the attending physician, when determining the individual regimen and dosage level at the most appropriate for a particular patient.

For preparing pharmaceutical compositions from the compounds of this invention, inert, pharmaceutically acceptable carriers can be either solid and liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories.

A solid carrier can be one or more substances, which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, or table disintegrating agents; it can also be an encapsulating material.

In powders, the carrier is a finely divided solid, which is in a mixture with the finely divided compound of the invention, or the active component. In tablets, the active component

- 14 -

is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

For preparing suppository compositions, a low-melting wax such as a mixture of fatty acid glycerides and cocoa butter is first melted and the active ingredient is dispersed therein by, for example, stirring. The molten homogeneous mixture is then poured into convenient sized moulds and allowed to cool and solidify.

Suitable carriers are magnesium carbonate, magnesium stearate, talc, lactose, sugar, pectin, dextrin, starch, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low-melting wax, cocoa butter, and the like.

The term composition is also intended to include the formulation of the active component with encapsulating material as a carrier providing a capsule in which the active component (with or without other carriers) is surrounded by a carrier which is thus in association with it. Similarly, cachets are included.

Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

Liquid form compositions include solutions, suspensions, and emulsions. For example, sterile water or water propylene glycol solutions of the active compounds may be liquid preparations suitable for parenteral administration. Liquid compositions can also be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions for oral administration can be prepared by dissolving the active component in water and adding suitable colorants, flavoring agents, stabilizers, and thickening agents as desired. Aqueous suspensions for oral use can be made by dispersing the finely divided active component in water together with a viscous material such as natural synthetic gums, resins, methyl cellulose, sodium carboxymethyl cellulose, and other suspending agents known to the pharmaceutical formulation art.

Depending on the mode of administration, the pharmaceutical composition will preferably include from 0.05% to 99%w (per cent by weight), more preferably from 0.10 to 50%w, of the compound of the invention, all percentages by weight being based on total composition.

A therapeutically effective amount for the practice of the present invention may be determined, by the use of known criteria including the age, weight and response of the individual patient, and interpreted within the context of the disease which is being treated or which is being prevented, by one of ordinary skills in the art.

- 15 -

Within the scope of the invention is the use of any compound of formula I as defined above for the manufacture of a medicament.

Also within the scope of the invention is the use of any compound of formula I for the manufacture of a medicament for the therapy of pain.

5 Additionally provided is the use of any compound according to Formula I for the manufacture of a medicament for the therapy of various pain conditions including, but not limited to: acute pain, chronic pain, neuropathic pain, back pain, cancer pain, and visceral pain.


10 A further aspect of the invention is a method for therapy of a subject suffering from any of the conditions discussed above, whereby an effective amount of a compound according to the formula I above, is administered to a patient in need of such therapy.

Additionally, there is provided a pharmaceutical composition comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable carrier.

15 Particularly, there is provided a pharmaceutical composition comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable carrier for therapy, more particularly for therapy of pain.

Further, there is provided a pharmaceutical composition comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, in association with a  
20 pharmaceutically acceptable carrier use in any of the conditions discussed above.


In another aspect, the present invention provides a method for preparing a compound of formula II,



25 comprising the steps of  
a) reacting a compound of formula III,




- 16 -

III

with a base having a pKa of more than 20;

b) reacting a product formed in step a) with a compound of formula IV,


IV

5 to form the compound of formula II,


wherein  $R^1$ ,  $R^2$ , and  $R^3$  are as previously defined, and  $R^c$  is  $C_{1-4}$ alkyl.

Particularly, the present invention provides a method of preparing a compound of formula II, wherein the strong base having a pKa of more than 20 is t-butyl lithium or n-butyl lithium.

10 In a further aspect, the present invention provides a process for preparing a compound of formula V,

V


comprising the step of reacting a compound of formula VI,

VI

15

with a compound of formula VII,

- 17 -


VII

to form the compound of formula V, wherein  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^e$  are defined as above and  $Y$  is CH or N.

Compounds of the present invention may be prepared according to the synthetic routes  
5 as depicted in Schemes 1 and 2 using one or more methods disclosed above.

- 18 -

Scheme 1



- 19 -

## Scheme 2

















Table 1 exemplifies some of the compounds of the present invention that were made according to the schemes and methods described above. These compounds were found to be active towards human CB1/CB2 receptors based on the test results of using one or more assays described below.

Table 1. Exemplary Compounds of the Invention.







Compound No.	Structure
1	
2	
3	
4	
5	
6	

- 21 -

Compound No.	Structure
7	
8	
9	
10	
11	
12	
13	









- 23 -







Compound No.	Structure
21	
22	
23	
24	
25	
26	



- 24 -

Compound No.	Structure
27	
28	
29	
30	
31	
32	

- 25 -

Compound No.	Structure
33	
34	
35	
36	
37	
38	

- 26 -

### Biological Evaluation

#### hCB<sub>1</sub> and hCB<sub>2</sub> receptor binding

Human CB<sub>1</sub> receptor from Receptor Biology (hCB<sub>1</sub>) or human CB<sub>2</sub> receptor from BioSignal (hCB<sub>2</sub>) membranes are thawed at 37 °C, passed 3 times through a 25-gauge blunt-end needle, diluted in the cannabinoid binding buffer (50 mM Tris, 2.5 mM EDTA, 5 mM MgCl<sub>2</sub>, and 0.5 mg/mL BSA fatty acid free, pH 7.4) and aliquots containing the appropriate amount of protein are distributed in 96-well plates. The IC<sub>50</sub> of the compounds of the invention at hCB<sub>1</sub> and hCB<sub>2</sub> are evaluated from 10-point dose-response curves done with <sup>3</sup>H-CP55,940 at 20000 to 25000 dpm per well (0.17-0.21 nM) in a final volume of 300 µl. The total and non-specific binding are determined in the absence and presence of 0.2 µM of HU210 respectively. The plates are vortexed and incubated for 60 minutes at room temperature, filtered through Unifilters GF/B (presoaked in 0.1% polyethyleneimine) with the Tomtec or Packard harvester using 3 mL of wash buffer (50 mM Tris, 5 mM MgCl<sub>2</sub>, 0.5 mg BSA pH 7.0). The filters are dried for 1 hour at 55 °C. The radioactivity (cpm) is counted in a TopCount (Packard) after adding 65 µl/well of MS-20 scintillation liquid.

Based on the above assays, the dissociation constant (K<sub>i</sub>) for a particular compound of the invention towards a particular receptor is determined using the following equation:

$$K_i = IC_{50} / (1 + [rad] / K_d),$$

Wherein IC<sub>50</sub> is the concentration of the compound of the invention at which 50% displacement has been observed;

[rad] is a standard or reference radioactive ligand concentration at that moment; and

K<sub>d</sub> is the dissociation constant of the radioactive ligand towards the particular receptor.

Using above-mentioned assays, the K<sub>i</sub> towards human CB<sub>1</sub> receptors for compounds 1-38 of the invention is measured to be in the range of 29 - 5852 nM. The K<sub>i</sub> towards human CB<sub>2</sub> receptors for compounds 1-38 of the invention is measured to be in the range of 0.7 - 753 nM.


### EXAMPLES

The invention will further be described in more detail by the following Examples which describe methods whereby compounds of the present invention may be prepared, purified, analyzed and biologically tested, and which are not to be construed as limiting the invention.


#### Example 1

- 27 -

*N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide:



Step A. *N*-(Cyclohexylmethyl)-3-methyl-5-nitro-2-pyridinamine:




5

To a solution of 2-chloro-3-methyl-5-nitropyridine (3.45 g, 20 mmol) in EtOH (100 mL) and triethylamine (5 mL) was added cyclohexylmethylamine (4.52 g, 40 mmol) at room temperature. The reaction mixture was refluxed for 12 hr, allowed to cool down to room temperature. After condensation, the residue was diluted with AcOEt, washed with 1 N


10 NH<sub>4</sub>OH and brine, dried over MgSO<sub>4</sub>. Removal of solvents provided the desired product (4.90 g, 98 %), which was used directly in the next step. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.02 (m, 2H), 1.23 (m, 3H), 1.75 (m, 6H), 2.16 (s, 3H), 3.46 (m, 2H), 4.98 (brs, 1H), 8.00 (s, 1H), 8.96 (s, 1H). MS (ESI) (M+H)<sup>+</sup> 250.31

15 Step B. *N*<sup>2</sup>-(cyclohexylmethyl)-3-methyl-2,5-pyridinediamine:




The above product was hydrogenated in ethyl acetate (150 mL) catalyzed by 10% Pd/C (200 mg) at 35-50 psi H<sub>2</sub> for 4 hr. The reaction mixture was filtered through Diatomaceous earth, and removal of solvents gave a product, which was purified by

20 flashmaster to give the desired product (4.14 g, 96 %). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.01 (m, 2H), 1.24 (m, 3H), 1.60 (m, 1H), 1.73 (m, 3H), 1.84 (m, 2H), 2.07 (s, 3H), 3.24 (d, J = 6.8 Hz, 2H), 6.79 (s, 1H), 7.61 (s, 1H). MS (ESI) (M+H)<sup>+</sup> 220.26

Step C. *N*-[6-[(cyclohexylmethyl)amino]-5-methyl-3-pyridinyl]-2,2-dimethyl-propanamide:

To a stirred solution of *N*<sup>2</sup>-(cyclohexylmethyl)-3-methyl-2,5-pyridinediamine (4.14 g, 18.9 mmol), diisopropylethylamine (5 mL) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) was added dropwise trimethylacetyl chloride (2.4 g, 20 mmol) at -50 °C. The reaction was allowed to warm up to 0 °C and then condensed under vacuum, diluted with AcOEt (200 mL), washed with 1N NH<sub>4</sub>OH (100 mL), brine (50 mL), and dried over MgSO<sub>4</sub>. Removal of solvent afforded the product as a solid (5.66 g, 99 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt): δ 1.02 (m, 2H), 1.27 (s, 9H), 1.28 (m, 3H), 1.80 (m, 6H), 2.27 (s, 3H), 3.25 (d, J = 7.6 Hz, 2H), 7.83 (s, 1H), 8.29 (s, 1H). MS (ESI) (M+H)<sup>+</sup> 303.30

Step D. *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide:

15

To a solution of *N*-[6-[(cyclohexylmethyl)amino]-5-methyl-3-pyridinyl]-2,2-dimethyl-propanamide (606 mg, 2.0 mmol) in dry THF was added a solution of BuLi (2.0 M, 4.5 mL, 9.0 mmol) at -50 °C. The reaction mixture was warmed up to -20 °C and stirred for an additional 1h at the temperature prior to addition of a solution of methyl 4-ethoxybenzeneacetic acid ester (392 mg, 2.0 mmol) in 1 mL THF. After 30 min, The reaction mixture was quenched with aqueous NH<sub>4</sub>Cl solution, and extracted with ethyl acetate (2 × 60 mL). The combined organic layers were washed with brine and dried over MgSO<sub>4</sub>. Removal of solvents gave a product, which was purified by Falshmaster to give the desired product 14

20

- 29 -


(350 mg, 39 %):  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ ):  $\delta$  1.08 (m, 5H), 1.32 (s, 9H), 1.35 (t,  $J = 6.4$  Hz, 3H), 1.42 (m, 2H), 1.69 (m, 4H), 4.00 (q,  $J = 6.4$  Hz, 2H), 4.02 (d,  $J = 7.6$  Hz, 2H), 4.15 (s, 2H), 6.34 (s, 1H), 6.87 (d,  $J = 8.4$  Hz, 2H), 7.14 (d,  $J = 8.4$  Hz, 2H), 8.34 (s, 1H), 8.55 (s, 1H).

Anal. Calcd. for  $\text{C}_{28}\text{H}_{37}\text{N}_3\text{O}_2 + 0.50 \text{ H}_2\text{O}$ : C, 73.65; H, 8.39. Found: C, 73.76; H, 8.65; Exact

5 mass Calcd. For  $\text{C}_{28}\text{H}_{37}\text{N}_3\text{O}_2 + 1$ , 448.2964, found: 448.3017 ( $\text{M}^+ + 1$ ).

### Example 2

*N*-[1-(cyclohexylmethyl)-2-[(3-methoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide:




10 Following the procedure 1D in the Example 1, using 3-methoxyphenylacetic anhydride (314 mg, 1.0 mmol) and *N*-[6-[(cyclohexylmethyl)amino]-5-methyl-3-pyridinyl]-2,2-dimethyl-propanamide (303 mg, 1.0 mmol), provided the desired title compound (234 mg, 54 %):  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , TFA salt):  $\delta$  1.12 (m, 5H), 1.36 (s, 9H), 1.52 (m, 2H), 1.72 (m, 4H), 3.81 (s, 3H), 4.05 (d,  $J = 7.6$  Hz, 2H), 4.20 (s, 2H), 6.36 (s, 1H), 6.92 (d,  $J = 8.4$  Hz, 2H), 7.20 (d,  $J = 8.4$  Hz, 2H), 8.29 (s, 1H), 8.51 (s, 1H). Anal. Calcd. for  $\text{C}_{27}\text{H}_{35}\text{N}_3\text{O}_2 + 0.70 \text{ TFA}$ : C, 66.44; H, 7.01, N, 8.18. Found: C, 66.23; H, 7.32, N, 7.84; MS (ESI) ( $\text{M} + \text{H}$ ) $^+$  434.02(MH $^+$ ).

### Example 3


*N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea

20



Step A. 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine

- 30 -



A solution of *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-  
 b]pyridin-5-yl]-2,2-dimethyl-propanamide (350 mg, 0.78 mmol) in dioxane (30 mL) and 20  
 % H<sub>2</sub>SO<sub>4</sub> (30 mL) was refluxed overnight, and then allowed to cool down to room  
 5 temperature. After condensation, the residue was diluted with AcOEt, washed with 1 N  
 NH<sub>4</sub>OH and brine, dried over MgSO<sub>4</sub>. Removal of solvents provided the desired product for  
 the next step (180 mg, 64 %), which was used directly in the next step. MS (ESI) (M+H)<sup>+</sup>  
 364.23.


To a stirred solution of the product formed in the last step (180 mg, 0.50 mmol),  
 10 diisopropylethylamine (1 mL) in CH<sub>2</sub>Cl<sub>2</sub> (30 mL) was added dropwise methyl chloroformate  
 (0.2 mL) at -30 °C. The reaction mixture was allowed to warm up to 0 °C, and then  
 condensed under vacuum. The residue was diluted with AcOEt, washed with 1 N NH<sub>4</sub>OH and  
 brine, dried over MgSO<sub>4</sub>. Removal of solvents provided a desired product, which was used  
 directly in the next step. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.86 -1.08 (m, 5H), 1.40 (t, J = 6.4 Hz, 3H),  
 15 1.63 (m, 6H), 3.75 (s, 3H), 3.98 (d, J = 7.6 Hz, 2H), 4.00 (q, J = 6.4 Hz, 2H), 4.02 (s, 2H),  
 6.06 (s, 1H), 6.82 (d, J = 8.4 Hz, 2H), 7.08 (d, J = 8.4 Hz, 2H), 8.00 (brs, 1H), 8.12 (s, 1H).  
 Exact mass Calcd. For C<sub>24</sub>H<sub>31</sub>N<sub>3</sub>O<sub>3</sub>+1, 422.2444, found: 422.2592 (M<sup>+</sup>+1).

To a solution of the product formed in the last step, methyl carbamate in THF was  
 dropwise added a solution of HCl (1M, 1 mL in diethyl ether) at -20 °C. After 10 min, LiAlH<sub>4</sub>  
 20 (0.72 g) was added to the solution. The reaction mixture was stirred at room temperature  
 overnight, and quenched carefully at -20 °C by adding MeOH (5 mL) and H<sub>2</sub>O (3 mL),  
 diluted with Et<sub>2</sub>O (50 mL), and then added Na<sub>2</sub>SO<sub>4</sub> (10 g). The resulting mixture was stirred  
 for 2 hr at r.t.. After filtration, the organic solution was concentrated *in vacuo* to afford a  
 product 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-  
 25 b]pyridin-5-amine (175 mg, 93 % for two steps), which was used in the next steps without  
 further purification. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.02 (m, 2H), 1.20 (m, 3H), 1.39 (t, J = 6.8 Hz, 3H),  
 1.54 (m, 2H), 1.64 (m, 3H), 1.82 (m, 1H), 2.84 (s, 3H), 3.92 (d, J = 7.6 Hz, 2H), 3.99 (q, J =

- 31 -

6.8 Hz, 2H), 4.03 (s, 2H), 5.96 (s, 1H), 6.83 (d,  $J = 8.4$  Hz, 2H), 7.05 (d,  $J = 2.4$  Hz, 1H), 7.10 (d,  $J = 8.4$  Hz, 2H), 7.80 (d,  $J = 2.4$  Hz, 1H). MS (ESI)  $(M+H)^+$  378.25.

Step B. *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea




A solution of 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (400 mg, 1.06 mmol), isopropyl isocyanate (425 mg, 5 mmol) and  $i\text{-Pr}_2\text{NEt}$  (1.0 mL) in  $\text{ClCH}_2\text{CH}_2\text{Cl}$  (30 mL) was refluxed for 1 h, and then concentrated.

- 10 The resulting residue was purified by preparative HPLC to give its TFA salt (204 mg, 33 %).  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , TFA salt):  $\delta$  1.03 (d,  $J = 6.4$  Hz, 6H), 1.08 (m, 5H), 1.34 (t,  $J = 6.8$  Hz, 3H), 1.46 (m, 2H), 1.64 (m, 3H), 1.76 (m, 1H), 3.23 (s, 3H), 3.88 (m, 1H), 3.98 (q,  $J = 7.0$  Hz, 2H), 3.98 (d,  $J = 7.6$  Hz, 2H), 4.12 (s, 2H), 6.24 (s, 1H), 6.83 (d,  $J = 8.8$  Hz, 2H), 7.10 (d,  $J = 8.8$  Hz, 2H), 7.84 (d,  $J = 2.4$  Hz, 1H), 8.06 (d,  $J = 2.4$  Hz, 1H). Exact mass Calcd. For  $\text{C}_{27}\text{H}_{38}\text{N}_4\text{O}_2+1$ , 463.3073, found: 463.3055 ( $M^++1$ ).

Example 4

*N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide



- 20 To a solution of 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (30 mg, 0.08 mmol) and  $i\text{-Pr}_2\text{NEt}$  (0.5 mL) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was added isovaleryl chloride (24 mg, 0.2 mmol) at 0 °C. The reaction mixture was stirred at room temperature overnight, and then concentrated. The resulting residue was purified by preparative HPLC to give its TFA salt (20 mg, 43 %).  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , TFA salt):  $\delta$  0.78




- 32 -

(d,  $J = 6.8$  Hz, 6H), 1.07 (m, 5H), 1.34 (t,  $J = 6.8$  Hz, 3H), 1.46 (m, 2H), 1.62 (m, 3H), 1.78 (m, 1H), 1.94 (d,  $J = 6.8$  Hz, 2H), 2.00 (m, 1H), 3.26 (s, 3H), 3.98 (q,  $J = 7.0$  Hz, 2H), 4.00 (d,  $J = 7.6$  Hz, 2H), 4.12 (s, 2H), 6.21 (s, 1H), 6.83 (d,  $J = 8.8$  Hz, 2H), 7.11 (d,  $J = 8.8$  Hz, 2H), 7.775 (d,  $J = 2.4$  Hz, 1H), 8.03 (d,  $J = 2.4$  Hz, 1H). MS (ESI)  $(M+H)^+$  462.07(MH<sup>+</sup>).

### 5 Example 5

*N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2-dimethyl-propanamide




Following the procedure in Example 4, using isobutyryl chloride (21 mg, 0.2 mmol) and 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (30 mg, 0.08 mmol), provided the desired compound as its TFA salt (22 mg, 49 %).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt):  $\delta$  0.98 (d,  $J = 6.8$  Hz, 6H), 1.10 (m, 5H), 1.35 (t,  $J = 6.8$  Hz, 3H), 1.48 (m, 2H), 1.64 (m, 3H), 1.78 (m, 1H), 2.46 (m, 1H), 3.26 (s, 3H), 3.99 (q,  $J = 7.0$  Hz, 2H), 4.00 (d,  $J = 7.6$  Hz, 2H), 4.13 (s, 2H), 6.24 (s, 1H), 6.84 (d,  $J = 8.8$  Hz, 2H), 7.11 (d,  $J = 8.8$  Hz, 2H), 7.85 (d,  $J = 2.4$  Hz, 1H), 8.09 (d,  $J = 2.4$  Hz, 1H). Exact mass Calcd. For C<sub>27</sub>H<sub>37</sub>N<sub>3</sub>O<sub>2</sub>+1, 448.2964, found: 448.3062 ( $M^+$ +1).

### Example 6

*N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-cyclopropanecarboxamide



20


Following the procedure in Example 4, using cyclopropanecarbonyl chloride (21 mg, 0.2 mmol) and 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (30 mg, 0.08 mmol), provided the desired compound as its TFA salt (24 mg, 54 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt):  $\delta$  0.63 (m, 2H), 0.90 (m, 2H), 1.08 (m, 6H), 1.35 (t,

- 33 -

J = 6.8 Hz, 3H), 1.48 (m, 2H), 1.64 (m, 3H), 1.78 (m, 1H), 3.29 (s, 3H), 3.99 (q, J = 7.0 Hz, 2H), 4.00 (d, J = 7.6 Hz, 2H), 4.11 (s, 2H), 6.24 (s, 1H), 6.84 (d, J = 8.8 Hz, 2H), 7.11 (d, J = 8.8 Hz, 2H), 7.90 (d, J = 2.4 Hz, 1H), 8.15 (d, J = 2.4 Hz, 1H). Exact mass Calcd. For  $C_{27}H_{35}N_3O_2+1$ , 446.2808, found: 446.2904 ( $M^++1$ ).

#### 5 Example 7


N-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-N,2,2-trimethyl-propanamide



Following the procedure in Example 4, using trimethylacetyl chloride (24 mg, 0.2 mmol) and 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-N-methyl-1H-pyrrolo[2,3-b]pyridin-5-amine (30 mg, 0.08 mmol), provided the desired compound as its TFA salt (25 mg, 54 %).  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , TFA salt):  $\delta$  1.04 (s, 9H), 1.08 (m, 5H), 1.35 (t, J = 7.0 Hz, 3H), 1.44 (m, 2H), 1.63 (m, 3H), 1.76 (m, 1H), 3.27 (s, 3H), 3.99 (q, J = 7.0 Hz, 2H), 4.02 (d, J = 7.6 Hz, 2H), 4.13 (s, 2H), 6.25 (s, 1H), 6.85 (d, J = 8.8 Hz, 2H), 7.13 (d, J = 8.8 Hz, 2H), 7.89 (d, J = 2.4 Hz, 1H), 8.11 (d, J = 2.4 Hz, 1H). Exact mass Calcd. For  $C_{24}H_{39}N_3O_2+1$ , 462.3121, found: 462.3208 ( $M^++1$ ).

#### Example 8

N-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-N',N'-diethyl-N-methyl-urea



20


Following the procedure in Example 4, using diethylcarbamyl chloride (27 mg, 0.2 mmol) and 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-N-methyl-1H-pyrrolo[2,3-b]pyridin-5-amine (30 mg, 0.08 mmol), provided the desired compound as its TFA salt (28 mg, 59 %).  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , TFA salt):  $\delta$  0.83 (t, J = 7.0 Hz, 6H), 1.09 (m, 5H), 1.35 (t, J =

- 34 -

7.0 Hz, 3H), 1.42 (m, 2H), 1.62 (m, 3H), 1.75 (m, 1H), 3.15 (q, J = 7.0 Hz, 4H), 3.15 (s, 3H), 3.99 (q, J = 7.0 Hz, 2H), 4.00 (d, J = 7.6 Hz, 2H), 4.11 (s, 2H), 6.22 (s, 1H), 6.84 (d, J = 8.8 Hz, 2H), 7.11 (d, J = 8.8 Hz, 2H), 7.82 (d, J = 2.4 Hz, 1H), 8.00 (d, J = 2.4 Hz, 1H). MS (ESI) (M+H)<sup>+</sup> 477.2

#### 5 Example 9


N-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-N,5-dimethyl-3-isoxazolecarboxamide



Following the procedure in Example 4, using 5-methyl-3-isoxazolecarbonyl chloride (50 mg, 0.33 mmol) and 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-N-methyl-1H-pyrrolo[2,3-b]pyridin-5-amine (60 mg, 0.16 mmol), provided the desired compound as its TFA salt (20 mg, 21 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt): δ 1.03 (m, 5H), 1.33 (t, J = 6.8 Hz, 3H), 1.38 (m, 2H), 1.62 (m, 4H), 2.20 (s, 3H), 3.45 (s, 3H), 3.94 (q, J = 7.0 Hz, 2H), 3.97 (d, J = 7.6 Hz, 2H), 4.07 (s, 2H), 5.93 (s, 1H), 6.12 (s, 1H), 6.82 (d, J = 8.4 Hz, 2H), 7.09 (d, J = 8.4 Hz, 2H), 7.75 (d, J = 2.0 Hz, 1H), 7.95 (d, J = 2.0 Hz, 1H). Anal. Calcd. for C<sub>29</sub>H<sub>34</sub>N<sub>4</sub>O<sub>3</sub> + 0.50 TFA: C, 66.28; H, 6.40; N, 10.31. Found: C, 66.24; H, 6.34; N, 10.22; Exact mass Calcd. For C<sub>29</sub>H<sub>34</sub>N<sub>4</sub>O<sub>3</sub>+1, 487.2709, found: 487.2712 (M<sup>+</sup>+1).

#### Example 10

N-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-2-fluoro-N-methyl-benzamide




Following the procedure in Example 4, using 2-fluorobenzoyl chloride (50 mg, 0.31 mmol) and 1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-N-methyl-1H-pyrrolo[2,3-

- 35 -


*b*]pyridin-5-amine (60 mg, 0.16 mmol), provided the desired compound as its TFA salt (30 mg, 31 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt): δ 0.88 ((m, 2H), 1.04 (m, 3H), 1.30 (m, 2H), 1.34 (t, J = 6.8 Hz, 3H), 1.60 (m, 4H), 3.48 (s, 3H), 3.87 (d, J = 8.0 Hz, 2H), 3.97 (q, J = 7.2 Hz, 2H), 4.03 (s, 2H), 6.04 (s, 1H), 6.82 (m, 3H), 6.99 (m, 1H), 7.06 (d, J = 8.4 Hz, 2H), 7.18 (m, 1H), 7.26 (m, 1H), 7.70 (d, J = 1.2 Hz, 1H), 7.86 (d, J = 1.2 Hz, 1H). Anal. Calcd. for C<sub>31</sub>H<sub>34</sub>FN<sub>3</sub>O<sub>2</sub> + 0.10 TFA: C, 73.33; H, 6.73; N, 8.22. Found: C, 72.93; H, 6.71; N, 8.19; Exact mass Calcd. For C<sub>31</sub>H<sub>34</sub>FN<sub>3</sub>O<sub>2</sub>+1, 500.2713, found: 500.2757 (M<sup>+</sup>+1).

Example 11

*N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide




Step A. 2-[(cyclohexylmethyl)amino]-5-[(2,2-dimethyl-1-oxopropyl)amino]-3-pyridineacetic acid:



To a solution of *N*-[6-[(cyclohexylmethyl)amino]-5-methyl-3-pyridinyl]-2,2-dimethyl-propanamide (303 mg, 1.0 mmol) in dry THF (20 mL) was added *t*-butyllithium (3.0 mL, 1.7 M, 5.1 mmol) at -50 °C. After warming up to -10 °C, CO<sub>2</sub> was introduced into the reaction mixture. After 10 min, the reaction mixture was quenched with aqueous NH<sub>4</sub>Cl solution, and extracted with ethyl acetate (2 × 100 mL). The combined organic layers were washed with brine and dried over MgSO<sub>4</sub>. Removal of solvents gave a product, which was consistent with the MS of the desired compound. The product was subject to the next step directly without purification.


- 36 -

Step B. Methyl-2-[(cyclohexylmethyl)amino]-5-[(2,2-dimethyl-1-oxopropyl)amino]-3-pyridineacetate:



To a solution of the product 2-[(cyclohexylmethyl)amino]-5-[(2,2-dimethyl-1-oxopropyl)amino]-3-pyridineacetic acid in dry MeOH (7.5 mL) was added 4N HCl solution (in dioxane, 2.5 mL) at 0 °C. The reaction mixture was stirred overnight at r.t, and then condensed under vacuum, diluted with AcOEt (50 mL), washed with 1N NH<sub>4</sub>OH (100 mL), brine (50 mL), and dried over MgSO<sub>4</sub>. Removal of solvent afforded desired title product (325 mg, 90 %). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.98 (m, 2H), 1.19 (m, 3H), 1.24 (s, 9H), 1.58 (m, 1H), 1.67 (m, 3H), 1.76 (m, 2H), 3.20 (d, J = 6.8 Hz, 2H), 3.39 (s, 2H), 3.63 (s, 3H), 4.98 (brs, 1H), 7.24 (brs, 1H), 7.680 (s, 1H), 7.92 (s, 1H).

Step C. *N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide



15 Intermediate 1


To a solution of 5-ethoxy-2-methyl-pyridine (274 mg, 2.0 mmol) in dry THF was added a solution of *t*-BuLi (1.7 M, 1.2mL, 2.04 mmol) at -78 °C. After stirring for about 3 min., a solution of Intermediate 1 (180 mg, 0.5 mmol) in 1.5 mL THF was added into the reaction mixture at -78 °C. The resulting mixture was stirred for an additional 30 min, and then quenched with aqueous NH<sub>4</sub>Cl solution, and extracted with ethyl acetate (2 × 50 mL). The combined organic layers were washed with brine and dried over MgSO<sub>4</sub>. Removal of solvents gave a product, which was purified by Gilson to give the desired product as its TFA

- 37 -


salts (95 mg, 28 %): <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt): δ 1.12 (m, 5H), 1.31 (s, 9H), 1.48 (t, J = 7.2 Hz, 3H), 1.52 (m, 2H), 1.70 (m, 3H), 1.82 (m, 1H), 4.08 (d, J = 7.6 Hz, 2H), 4.25 (q, J = 6.8 Hz, 2H), 4.53 (s, 2H), 6.13 (s, 1H), 7.72 (d, J = 8.8 Hz, 1H), 7.99 (dd, J = 8.8, 2.8 Hz, 1H), 8.08 (d, J = 2.4 Hz, 1H), 8.30 (d, J = 2.4 Hz, 1H), 8.45 (d, J = 2.8 Hz, 1H). MS (ESI) (M+H)<sup>+</sup> 449.2

Example 12

[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-, 1-methylethyl ester carbamic acid




10 Step A. 1-Methylethyl [6-[(cyclohexylmethyl)amino]-5-methyl-3-pyridinyl]-carbamic acid ester



Following the same method as described for preparing the compound in Example 1 step C, using 2.75 g (12.6 mmol) of *N*<sup>2</sup>-(cyclohexyl methyl)-3-methyl-2,5-pyridinediamine and isopropyl chloroformate (1 M in toluene, 13 ml, 13 mmol), provided the title compound (3.87 g, 100%). <sup>1</sup>H-NMR (CD<sub>3</sub>OD): δ 1.00 (m, 2H), 1.20 (m, 3H), 1.26 (d, J = 6.4 Hz, 6H), 1.58 (m, 1H), 1.72 (m, 3H), 1.80 (m, 2H), 2.06 (s, 3H), 3.24 (d, J = 6.8 Hz, 2H), 4.98 (m, 1H), 6.29 (brs, 1H), 7.50 (s, 1H), 7.83 (s, 1H). MS (ESI) (M+H)<sup>+</sup>: 305.30.


20 Step B. 2-[(cyclohexylmethyl)amino]-5-[[[(1-methylethoxy)carbonyl]amino]-3-pyridineacetic acid methyl ester

- 38 -



Following the same method as described for the synthesis of the compound in Example 11, step A, using starting material 1-methylethyl [6-[(cyclohexylmethyl)amino]-5-methyl-3-pyridinyl]-carbamic acid ester (1.22 g, 4.0 mmol, in dry THF 960 mL) and t-butyllithium  
 5 (9.5 mL, 1.7 M, 16.0 mmol), provided the title compound.


Step C. 2-[(cyclohexylmethyl)amino]-5-[[[(1-methylethoxy)carbonyl]amino]-3-pyridineacetic acid methyl ester



10 Following the same method as described for preparing methyl-2-  
 [(cyclohexylmethyl)amino]-5-[(2,2-dimethyl-1-oxopropyl)amino]-3-pyridineacetate  
 (Example 11, Step B). Using the product from the last step as the starting material, provided  
 a product, which was purified by Gilson followed by work up to give the product (765 mg, 53  
 15 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD): δ 1.00 (m, 2H), 1.20 (m, 3H), 1.26 (d, J = 6.4 Hz, 6H), 1.58 (m, 1H),  
 1.72 (m, 3H), 1.80 (m, 2H), 3.24 (d, J = 6.8 Hz, 2H), 3.44 (s, 2H), 3.67 (s, 3H), 4.05 (brs, 1H),  
 4.98 (m, 1H), 6.32 (brs, 1H), 7.60 (s, 1H), 7.90 (s, 1H).

Step D. [1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-, 1-methylethyl ester carbamic acid


- 39 -




- Method as described for *N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide (Example 11, Step C), but using 5-ethoxy-2-methyl-pyridine (1.37 g, 10.0 mmol) and 2-[(cyclohexylmethyl)amino]-5-[[1-methylethoxy]carbonyl]amino]-3-pyridineacetic acid methyl ester (726 mg, 2.0 mmol),
- 5 provided the title compound as its TFA salt (546 mg, 40 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt): δ 1.11 (m, 5H), 1.29 (d, *J* = 6.0 Hz, 6H), 1.46 (t, *J* = 7.00Hz, 3H), 1.50 (m, 2H), 1.72 (m, 3H), 1.81 (m, 1H), 4.08 (d, *J* = 8.0 Hz, 2H), 4.25 (q, *J* = 6.8Hz, 2H), 4.55 (s, 2H), 4.93 (m, 1H), 6.13 (s, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 8.02 (dd, *J* = 8.8, 2.8 Hz, 1H), 8.09 (s, 1H), 8.26 (s, 1H),
- 10 8.47 (d, *J* = 2.8 Hz, 1H). MS (ESI) (*M*+*H*)<sup>+</sup> 451.2.

**Example 13**

*N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2,2-trimethyl-propanamide



- 15 Step A. 1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine




To a solution of [1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-, 1-methylethyl ester carbamic acid



- 40 -

(520 mg, 0.77 mmol) in THF was added  $\text{LiAlH}_4$  (1.44 g) at  $-20^\circ\text{C}$ . The reaction mixture was stirred at room temperature overnight, and quenched carefully at  $-20^\circ\text{C}$  by adding MeOH (5 mL) and  $\text{H}_2\text{O}$  (3 mL), diluted with  $\text{Et}_2\text{O}$  (50 mL), and then added  $\text{Na}_2\text{SO}_4$  (10 g). The resulting mixture was stirred for 2 hr at r.t.. After filtration, the organic solution was concentrated *in vacuo* to afford a product (263 mg, 90 %), which was used in the next steps without further purification.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.08 (m, 5H), 1.35 (t,  $J = 7.2$  Hz, 3H), 1.42 (m, 2H), 1.62 (m, 3H), 1.75 (m, 1H), 3.15 (s, 3H), 3.99 (q,  $J = 7.2$  Hz, 2H), 4.00 (d,  $J = 7.2$  Hz, 2H), 4.11 (s, 2H), 6.22 (s, 1H), 6.84 (d,  $J = 8.4$  Hz, 1H), 7.11 (d,  $J = 8.4$  Hz, 1H), 7.60 (brs, 1H), 7.82 (s, 1H), 8.00 (s, 1H). MS (ESI)  $(\text{M}+\text{H})^+$  379.94.

Step B. *N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2,2-trimethyl-propanamide




Following the procedure in Example 7, using 1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine

(30 mg, 0.08 mmol) and trimethylacetyl chloride (24 mg, 0.2 mmol), provided the desired compound as its TFA salt (15 mg, 27 %).  $^1\text{H-NMR}$  ( $\text{CD}_3\text{OD}$ , TFA salt):  $\delta$  1.01 (s, 9H), 1.12 (m, 5H), 1.46 (t,  $J = 7.2$  Hz, 3H), 1.50 (m, 2H), 1.72 (m, 3H), 1.84 (m, 1H), 3.25 (s, 3H), 4.13 (d,  $J = 7.2$  Hz, 2H), 4.25 (q,  $J = 7.2$  Hz, 2H), 4.55 (s, 2H), 6.16 (s, 1H), 7.72 (d,  $J = 8.8$  Hz, 1H), 7.82 (d,  $J = 2.4$  Hz, 1H), 7.97 (dd,  $J = 8.8, 2.8$  Hz, 1H), 8.14 (d,  $J = 2.4$  Hz, 1H), 8.45 (d,  $J = 2.8$  Hz, 1H). MS (ESI)  $(\text{M}+\text{H})^+$  463.2

Example 14

*N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide


- 41 -



Following the procedure in Example 4, using 1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (30 mg, 0.08 mmol) and isobutyryl chloride (24 mg, 0.2 mmol), provided the desired compound as its TFA salt (12 mg, 22 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt): δ 0.79 (d, *J* = 6.4 Hz, 6H), 1.12 (m, 5H), 1.46 (t, *J* = 7.2 Hz, 3H), 1.54 (m, 2H), 1.68 (m, 3H), 1.84 (m, 1H), 1.93 (d, *J* = 6.8 Hz, 2H), 2.01 (m, 1H), 3.28 (s, 3H), 4.13 (d, *J* = 7.6 Hz, 2H), 4.25 (q, *J* = 6.8 Hz, 2H), 4.56 (s, 2H), 6.18 (s, 1H), 7.72 (d, *J* = 9.2 Hz, 1H), 7.80 (d, *J* = 2.4 Hz, 1H), 7.98 (dd, *J* = 9.2, 2.8 Hz, 1H), 8.12 (d, *J* = 2.4 Hz, 1H), 8.46 (d, *J* = 2.8 Hz, 1H). MS (ESI) (*M*+*H*)<sup>+</sup> 463.2

#### 10 Example 15

*N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea




Following the procedure B in Example 3, using 1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-*N*-methyl-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (30 mg, 0.08 mmol) and isopropyl isocyanate (43 mg, 0.5 mmol), provided the desired compound as its TFA salt (20 mg, 36 %). <sup>1</sup>H-NMR (CD<sub>3</sub>OD, TFA salt): δ 1.06 (d, *J* = 6.4 Hz, 6H), 1.12 (m, 5H), 1.46 (t, *J* = 7.2 Hz, 3H), 1.54 (m, 2H), 1.68 (m, 3H), 1.84 (m, 1H), 3.25 (s, 3H), 3.88 (m, 1H), 4.11 (d, *J* = 7.6 Hz, 2H), 4.25 (q, *J* = 7.2 Hz, 2H), 4.58 (s, 2H), 6.18 (s, 1H), 7.75 (d, *J* = 8.8 Hz, 1H), 7.83 (d, *J* = 2.4 Hz, 1H), 8.04 (dd, *J* = 8.8, 2.8 Hz, 1H), 8.14 (d, *J* = 2.4 Hz, 1H), 8.49 (d, *J* = 2.8 Hz, 1H).


#### Example 16

*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide

- 42 -



Step A. 2,2-dimethyl-N-[5-methyl-6-[[[(tetrahydro-2H-pyran-4-yl)methyl]amino]-3-pyridinyl]-propanamide



5 To a solution of 2-chloro-3-methyl-5-nitropyridine (5.0 g, 29.0 mmol) in ethanol (100 ml) at room temperature was added triethylamine (8.0 ml, 58.0 mmol) followed by 4-aminomethyl tetrahydropyran (3.7 g, 31.9 mmol). The reaction mixture was refluxed overnight. Subsequently, the mixture was cooled to room temperature and concentrated *in vacuo*.

10 The residue was taken up into ethyl acetate (75 ml) and palladium on carbon (120 mgs, 10% grade, 0.1 mmol) was added. The suspension was placed in Parr apparatus and shaken for 72 hours under a hydrogen atmosphere (35 psi). The suspension was then brought to normal atmosphere and filtered on Diatomaceous earth. The filtrate was concentrated *in vacuo*.


15 This residue was taken up into dichloromethane (125 ml) at -78°C to which was added diisopropyl ethylamine (6.1 ml, 34.8 mmol) followed by pivaloyl chloride (3.73 ml, 30.3 mmol). The mixture was stirred for two hours at 0°C and then quenched with 2M NaOH aqueous solution (50 ml). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (125 ml). The organic phases were combined, dried  
20 with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using silica gel flash chromatography ([3% MeOH + 0.5% NH<sub>4</sub>OH<sub>aq</sub>] in CH<sub>2</sub>Cl<sub>2</sub>) to provide 8.1 g of the title compound 2,2-dimethyl-N-[5-methyl-6-[[[(tetrahydro-2H-pyran-4-yl)methyl]amino]-3-

- 43 -

pyridinyl]-propanamide. <sup>1</sup>H NMR (400 MHz, CHLOROFORM-D)  $\delta$  ppm 1.29 (s, 9 H) 1.36 (m, 2 H) 1.67 (m, 2 H) 1.88 (m, 1 H) 2.04 (d,  $J=10.55$  Hz, 3 H) 3.36 (m, 4 H) 3.97 (m, 2 H) 4.12 (m, 1 H) 7.12 (s, 1 H) 7.63 (d,  $J=1.76$  Hz, 1 H) 7.86 (d,  $J=2.73$  Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 306.

5

Step B. *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide




To a solution of 2,2-dimethyl-*N*-[5-methyl-6-[(tetrahydro-2*H*-pyran-4-yl)methyl]amino]-3-pyridinyl]-propanamide (2.0 g, 6.55 mmol) in THF (70 ml) at -78°C was added *n*-butyl lithium (11.5 ml of 2.0 M solution in cyclohexane, 23.0 mmol). The mixture was stirred for one hour at -20°C and then cooled to -78°C. To this reaction mixture was cannulated a solution of methyl 4-ethoxybenzoate (1.72g, 7.88 mmol) in THF (50 ml) at -78°C. After stirring for 3 hours at room temperature, the mixture was quenched with NaHCO<sub>3</sub> saturated aqueous solution (200 ml) and diluted with EtOAc (100 ml). The phases were separated and the aqueous phase was back-extracted with additional EtOAc (100 ml). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using silica gel flash chromatography ([3% MeOH + 0.5% NH<sub>4</sub>OHaq] in CH<sub>2</sub>Cl<sub>2</sub>) to provide 750 mg of the title compound *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide. <sup>1</sup>H NMR (400 MHz, CHLOROFORM-D)  $\delta$  ppm 1.31 (m, 2 H) 1.34 (s, 9 H) 1.40 (t,  $J=7.03$  Hz, 3 H) 1.56 (m, 2 H) 2.04 (m, 1 H) 3.22 (m, 2 H) 3.35 (m, 2 H) 3.90 (t,  $J=3.32$  Hz, 2 H) 4.00 (m, 2 H) 4.06 (s, 2 H) 6.12 (s, 1 H) 6.83 (m, 2 H) 7.07 (d,  $J=8.79$  Hz, 2 H) 7.32 (m, 1 H) 8.09 (d,  $J=2.34$  Hz, 1 H) 8.15 (d,  $J=2.34$  Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 451.


Example 17:

- 44 -

*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,6-difluoro-*N*-methyl-benzenesulfonamide



Step A. [2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]- carbamic acid methyl ester



*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide (1.3 g, 3.0 mmol) was dissolved into a mixture of dioxane (25 mL) and 20% sulfuric acid aqueous solution (25 mL) at room temperature. The solution was brought to 120°C. After stirring for 6 hours, the mixture was cooled to room temperature and concentrated *in vacuo*. The residue was brought to pH 8 by addition of 2M NaOH aqueous solution (200 mL). The mixture was extracted twice with EtOAc (100 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*.


The residue was taken up into dichloromethane (50 mL) and the mixture cooled to -30°C. To this solution was added diisopropyl ethylamine (1.1 mL, 6.2 mmol) followed by a solution of methylchloroformate (231 µL, 3.0 mmol) in dichloromethane (25 mL) at -78°C. The reaction was allowed to warm to 0°C. After stirring for 3 hours, the reaction was quenched with 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (50 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (50 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using silica gel flash chromatography ([3% MeOH + 0.5% NH<sub>4</sub>OHaq] in CH<sub>2</sub>Cl<sub>2</sub>) to provide 374 mg of the title compound [2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-carbamic acid methyl

- 45 -

ester.  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $\text{D}$ )  $\delta$  1.29 (m, 2 H), 1.39 (t,  $J=6.93$  Hz, 3 H), 1.56 (m, 2 H), 2.06 (m, 1 H), 3.35 (td,  $J=11.81, 2.15$  Hz, 2 H), 3.77 (s, 3 H), 3.89 (m, 2 H), 3.95 (m, 2 H), 3.99 (m, 2 H), 4.05 (s, 2 H), 5.72 (s, 1 H), 6.09 (s, 1 H), 6.83 (m, 2 H), 7.07 (d,  $J=8.79$  Hz, 2 H), 8.08 (d,  $J=2.34$  Hz, 1 H), 8.14 (d,  $J=2.34$  Hz, 1 H). MS (ESI)  $(\text{M}+\text{H})^+$ : 425.

5


Step B. 2-[(4-ethoxyphenyl)methyl]-N-methyl-1-[(tetrahydro-2H-pyran-4-yl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-amine



To a solution of [2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2H-pyran-4-yl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-carbamic acid methyl ester (153 mg, 0.36 mmol) in THF (25 ml) at  $0^\circ\text{C}$  was added lithium aluminum hydride (35 mg, 0.90 mmol). The mixture was stirred for 48 hours at room temperature. The reaction was then cooled to  $0^\circ\text{C}$  and quenched by dropwise addition of water (35  $\mu\text{L}$ ), followed by 4M NaOH aqueous solution (35  $\mu\text{L}$ ) and water (105  $\mu\text{L}$ ). After stirring at room temperature for 30 minutes, the suspension was filtered over Diatomaceous earth and concentrated *in vacuo* to give 100 mgs of colorless oil.  $^1\text{H}$  NMR (400 MHz, CHLOROFORM- $\text{D}$ )  $\delta$  1.40 (m, 5 H), 1.59 (m, 2 H), 2.05 (m, 1 H), 2.87 (s, 3 H), 3.36 (m, 2 H), 3.66 (m, 2 H), 4.01 (m, 6 H), 5.99 (s, 1 H), 6.83 (d,  $J=8.79$  Hz, 2 H), 6.98 (s, 1 H), 7.05 (d,  $J=2.54$  Hz, 1 H), 7.10 (d,  $J=8.79$  Hz, 2 H), 7.79 (d,  $J=2.54$  Hz, 1 H). MS (ESI)  $(\text{M}+\text{H})^+$ : 380.

20

Step C. N-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2H-pyran-4-yl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-2,6-difluoro-N-methyl-benzenesulfonamide




- 46 -

To a solution of 2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (51 mg, 0.13 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (68 µL, 0.39 mmol) followed by 2,6-difluorobenzenesulfonyl chloride (57 mg, 0.27 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2 M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 36 mg of the TFA salt of *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,6-difluoro-*N*-methyl-benzenesulfonamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 1.20 (m, 4 H), 1.26 (t, *J*=7.03 Hz, 3 H), 1.82 (m, 1 H), 3.09 (m, 2 H), 3.31 (s, 3 H), 3.73 (m, 2 H), 3.90 (m, 2 H), 3.94 (d, *J*=7.81 Hz, 2 H), 4.02 (s, 2 H), 6.04 (s, 1 H), 6.75 (d, *J*=8.79 Hz, 2 H), 7.01 (m, 4 H), 7.55 (m, 2 H), 7.89 (d, *J*=2.34 Hz, 1 H). MS (ESI) (M+H)<sup>+</sup> : 557.

#### Example 18

*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-cyclobutanecarboxamide




To a solution of 2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (54 mg, 0.14 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (74 µL, 0.42 mmol) followed by cyclobutanecarbonyl chloride (33 µL, 0.28 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 25 mg of the TFA salt of *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-

- 47 -

*N*-methyl-cyclobutanecarboxamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 1.26 (m, 5 H), 1.61 (m, 4 H), 1.89 (m, 1 H), 2.13 (m, 2 H), 2.95 (m, 2 H), 3.11 (m, 2 H), 3.16 (s, 3 H), 3.75 (m, 2 H), 3.91 (q, *J*=6.96 Hz, 2 H), 3.98 (d, *J*=7.42 Hz, 2 H), 4.04 (m, 1 H), 4.06 (s, 2 H), 6.14 (s, 1 H), 6.77 (d, *J*=8.59 Hz, 2 H), 7.05 (d, *J*=8.59 Hz, 2 H), 7.62 (d, *J*=2.34 Hz, 1 H), 7.90 (d, *J*=2.34 Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 463.

**Example 19**

*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,5-difluoro-*N*-methyl-benzamide




To a solution of 2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (52 mg, 0.14 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (70 μL, 0.41 mmol) followed by 2,5-difluorophenylcarbonyl chloride (35 μL, 0.27 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2 M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 28 mg of the TFA salt of *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,5-difluoro-*N*-methyl-benzamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 1.17 (m, 4 H), 1.26 (t, *J*=6.93 Hz, 3 H), 1.75 (m, 1 H), 3.05 (m, 2 H), 3.40 (s, 3 H), 3.69 (m, 2 H), 3.90 (m, 4 H), 3.99 (s, 2 H), 6.03 (s, 1 H), 6.75 (d, *J*=8.59 Hz, 2 H), 6.78 (m, 1 H), 6.83 (m, 1 H), 7.01 (d, *J*=8.59 Hz, 2 H), 7.04 (m, 1 H), 7.66 (d, *J*=2.15 Hz, 1 H), 7.85 (d, *J*=2.15 Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 521.

**Example 20**

*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2-dimethyl-propanamide




- 48 -



To a solution of 2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (50 mg, 0.13 mmol) in dichloromethane (5 ml) at 0°C was added diisopropylethylamine (68 µL, 0.39 mmol) followed by isobutanoyl chloride (28 µL, 0.26 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 28 mg of the TFA salt of *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2-dimethyl-propanamide. <sup>1</sup>H NMR (400 MHz, METHANOL-*D*<sub>4</sub>) δ 0.71 (d, *J*=6.64 Hz, 6 H) 1.19 (m, 2H) 1.26 (m, 5 H) 1.90 (m, 1 H) 2.39 (m, 1 H) 3.11 (m, 2H) 3.17 (s, 3 H) 3.76 (m, 2 H) 3.91 (m, 2 H) 3.98 (d, *J*=7.42 Hz, 2 H) 4.06 (s, 2 H) 6.16 (s, 1 H) 6.77 (d, *J*=7.82 Hz, 2 H) 7.05 (d, *J*=7.82 Hz, 2 H) 7.71 (s, 1 H) 7.98 (s, 1 H). MS (ESI) (M+H)<sup>+</sup> : 451.

#### Example 21

*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2,2-trimethyl-propanamide




To a solution of 2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (53 mg, 0.14 mmol) in dichloromethane (5 ml) at 0°C was added diisopropylethylamine (73 µL, 0.42 mmol) followed by 2,2-dimethylpropanoyl chloride (34 µL, 0.28 mmol). The mixture was stirred overnight at room

- 49 -

temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 20 mg of the TFA salt of *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2,2-trimethyl-propanamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 0.92 (s, 9 H) 1.25 (m, 7 H) 1.87 (m, 1 H) 3.10 (m, 2 H) 3.15 (s, 3 H) 3.75 (m, 2 H) 3.91 (q, *J*=6.96 Hz, 2 H) 3.99 (d, *J*=7.42 Hz, 2 H) 4.06 (s, 2 H) 6.14 (s, 1 H) 6.77 (d, *J*=8.79 Hz, 2 H) 7.06 (d, *J*=8.79 Hz, 2 H) 7.70 (d, *J*=2.34 Hz, 1 H) 7.97 (d, *J*=2.34 Hz, 1 H). MS (ESI) (*M*+*H*)<sup>+</sup>: 465

**Example 22**

*N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea




To a solution of 2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (710 mg, 1.87 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (977 μL, 5.61 mmol) followed by isopropylisocyanate (367 μL, 3.74 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (100 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (100 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 461 mg of the TFA salt of *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea. <sup>1</sup>H NMR (400 MHz, CHLOROFORM-D) δ 1.02 (d, *J*=6.44 Hz, 6 H) 1.40 (t, *J*=6.93 Hz, 3 H) 1.45 (m, 4 H) 2.13 (m, 2 H) 3.26 (m, 2 H) 3.27 (s, 3 H) 3.94 (m, 2 H) 4.01 (q, *J*=6.93 Hz, 2 H) 4.05 (d, *J*=6.93 Hz, 2 H) 4.09 (s, 2 H) 6.14 (s, 1 H)

- 50 -

6.86 (m, 3 H) 7.10 (d,  $J=8.59$  Hz, 2 H) 7.65 (d,  $J=2.34$  Hz, 1 H) 8.13 (d,  $J=2.34$  Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 466

Example 23

5 *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide




To a solution of 2-[(4-ethoxyphenyl)methyl]-*N*-methyl-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (68 mg, 0.18 mmol) in dichloromethane (5 ml) at 0°C was added diisopropylethylamine (94 µL, 0.54 mmol) followed by isovaleryl chloride (44 µL, 0.36 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using reversed phase silica gel flash chromatography (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 33 mg of the TFA salt of *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 0.71 (d,  $J=6.44$  Hz, 6 H) 1.26 (t,  $J=6.93$  Hz, 3 H) 1.33 (m, 4 H) 1.85 (m, 2 H) 1.89 (m, 2 H) 3.20 (s, 3 H) 3.74 (m, 2 H) 3.90 (q,  $J=6.93$  Hz, 2 H) 3.98 (d,  $J=7.23$  Hz, 2 H) 4.04 (m, 2 H) 4.06 (s, 2 H) 6.14 (s, 1 H) 6.76 (d,  $J=8.59$  Hz, 2 H) 7.05 (d,  $J=8.59$  Hz, 2 H) 7.66 (d,  $J=2.34$  Hz, 1 H) 7.93 (d,  $J=2.34$  Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 465.


Example 24

*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide

- 51 -



Step A: *N*-[6-[(cyclobutylmethyl)amino]-5-methyl-3-pyridinyl]-2,2-dimethyl-propanamide




To a solution of 2-chloro-3-methyl-5-nitropyridine (5.1 g, 29.7 mmol) in ethanol (100 mL) at room temperature was added triethylamine (8.3 ml, 59.4 mmol) followed by cyclobutyl methylamine (2.8 g, 32.7 mmol). The reaction mixture was refluxed overnight. Subsequently, the mixture was cooled to room temperature and concentrated *in vacuo*.

The residue was taken up into ethyl acetate (75 ml) and palladium on carbon (120 mg, 10% grade, 0.1 mmol) was added. The suspension was placed in Parr apparatus and shaken for 72 hours under a hydrogen atmosphere (35 psi). The suspension was then brought to normal atmosphere and filtered on Diatomaceous earth. The filtrate was concentrated *in vacuo*.

This residue was taken up into dichloromethane (125 mL) at -78°C to which was added diisopropyl ethylamine (6.2 mL, 35.6 mmol) followed by pivaloyl chloride (3.84 ml, 31.2 mmol). The mixture was stirred for two hours at 0°C and then quenched with 2M NaOH aqueous solution (50 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (125 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using silica gel flash chromatography ([3% MeOH + 0.5% NH<sub>4</sub>OHaq] in CH<sub>2</sub>Cl<sub>2</sub>) to provide 6.75 g of the title compound *N*-[6-[(cyclobutylmethyl)amino]-5-methyl-3-pyridinyl]-2,2-dimethyl-propanamide. <sup>1</sup>H NMR (400 MHz, CHLOROFORM-D) δ 1.29 (s, 9 H) 1.74 (m, 2 H) 1.91 (m, 2 H) 2.05 (s, 3 H) 2.09 (m, 2 H) 2.57 (m, 1 H) 3.45 (dd, *J*=7.42, 5.27 Hz, 2 H) 3.96 (s, 1 H) 7.09 (s, 1 H) 7.65 (d, *J*=2.54 Hz, 1 H) 7.85 (d, *J*=2.54 Hz, 1 H). MS (ESI) (*M*+H)<sup>+</sup>: 276.

- 52 -


Step B. *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide



To a solution of *N*-[6-[(cyclobutylmethyl)amino]-5-methyl-3-pyridinyl]-2,2-dimethyl-propanamide (2.0 g, 7.26 mmol) in THF (70 ml) at -78°C was added *n*-butyl lithium (12.7 mL of 2.0 M solution in cyclohexane, 25.4 mmol). The mixture was stirred for one hour at -20°C and then cooled to -78°C. To this reaction mixture was cannulated to a solution of methyl 4-ethoxy-benzoate (2.3 g, 11.6 mmol) in THF (50 mL) at -78°C. After stirring for 3 hours at room temperature, the mixture was quenched with NaHCO<sub>3</sub> saturated aqueous solution (200 ml) and diluted with EtOAc (100 mL). The phases were separated and the aqueous phase was back-extracted with additional EtOAc (100 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using silica gel flash chromatography ([3% MeOH + 0.5% NH<sub>4</sub>OHaq] in CH<sub>2</sub>Cl<sub>2</sub>) to provide 650 mg of the title compound *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.21 (s, 9 H) 1.29 (t, *J*=7.03 Hz, 3 H) 1.77 (m, 6 H) 2.69 (m, 1 H) 3.97 (q, *J*=7.03 Hz, 2 H) 4.06 (s, 2 H) 4.15 (d, *J*=7.23 Hz, 2 H) 5.99 (s, 1 H) 6.86 (d, *J*=8.59 Hz, 2 H) 7.14 (d, *J*=8.59 Hz, 2 H) 7.98 (d, *J*=2.34 Hz, 1 H) 8.21 (d, *J*=2.34 Hz, 1 H) 9.21 (s, 1 H). MS (ESI) (*M*+*H*)<sup>+</sup>: 421.

Example 25

[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-, methyl ester carbamic acid




*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide (872.4 mg, 2.1 mmol) was dissolved into a mixture of dioxane (25 mL) and 20% sulfuric acid aqueous solution (25 mL) at room temperature. The solution was  
 5 brought to 120°C. After stirring for 6 hours, the mixture was cooled to room temperature and concentrated *in vacuo*. The residue was brought to pH 8 by addition of 2M NaOH aqueous solution (200 mL). The mixture was extracted twice with EtOAc (100 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*.

The residue was taken up into dichloromethane (50 mL) and the mixture cooled to –  
 10 30°C. To this solution was added diisopropyl ethylamine (905 µL, 5.2 mmol) followed by a solution of methylchloroformate (193 µL, 2.5 mmol) in dichloromethane (25 mL) at -78°C. The reaction was allowed to warm to 0°C. After stirring for 3 hours, the reaction was quenched with 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (50 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (50 mL). The organic  
 15 phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using silica gel flash chromatography ([3% MeOH + 0.5% NH<sub>4</sub>OHaq] in CH<sub>2</sub>Cl<sub>2</sub>) to provide 701 mg of the title compound [1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-methyl ester carbamic acid. <sup>1</sup>H NMR (400 MHz, CHLOROFORM-D) δ 1.37 (t, J=6.93 Hz, 3 H) 1.78 (m, 6 H) 2.66 (m, 1 H) 3.75 (s, 3 H) 3.89 (q, J=6.93 Hz, 2 H) 4.01 (s, 2 H) 4.11 (d, J=7.23 Hz, 2 H) 5.44 (s, 1 H) 6.08 (s, 1 H) 6.85 (m, 2 H) 7.09 (m, 2 H) 8.05 (d, J=2.35 Hz, 1 H) 8.13 (d, J=2.35 Hz, 1 H) MS (ESI) (M+H)<sup>+</sup> : 395.


#### Example 26

*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,6-  
 25 difluoro-*N*-methyl-benzenesulfonamide




Step A. 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-  
amine

- 54 -



To a solution of [1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-  
*b*]pyridin-5-yl]-methyl ester carbamic acid (701 mg, 1.78 mmol) in THF (25 mL) at 0°C was  
 added lithium aluminum hydride (300 mg, 7.90 mmol). The mixture was stirred for 48 hours  
 at room temperature. The reaction was then cooled to 0°C and quenched by dropwise addition  
 of water (300 µL), followed by 4M NaOH aqueous solution (300 µL) and water (900 µL).  
 After stirring at room temperature for 30 minutes, the suspension was filtered over  
 Diatomaceous earth and concentrated *in vacuo* to give 517 mgs of colorless oil. MS (ESI)  
 (M+H)<sup>+</sup>: 350.

Step B. *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-  
 yl]-2,6-difluoro-*N*-methyl-benzenesulfonamide




To a solution 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]- 1*H*-pyrrolo[2,3-  
*b*]pyridin-5-amine (50 mg, 0.14 mmol) in dichloromethane (5 mL) at 0°C was added  
 diisopropylethylamine (75 µL, 0.43 mmol) followed by 2,6-difluorobenzenesulfonyl chloride  
 (61 mg, 0.29 mmol). The mixture was stirred overnight at room temperature. The reaction  
 was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and  
 the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic  
 phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue  
 was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous  
 solution) to provide 24 mg of the TFA salt of *N*-[1-(cyclobutylmethyl)-2-[(4-  
 ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,6-difluoro-*N*-methyl-  
 benzenesulfonamide. <sup>1</sup>H NMR (400 MHz, METHANOL-*D*<sub>4</sub>) δ 1.35 (t, *J*=7.03 Hz, 3 H) 1.82

- 55 -

(m, 6 H) 2.71 (m, 1 H) 3.40 (s, 3 H) 3.99 (q,  $J=7.03$  Hz, 2 H) 4.08 (s, 2 H) 4.17 (d,  $J=7.23$  Hz, 2 H) 6.05 (s, 1 H) 6.84 (m, 2 H) 7.09 (m, 4 H) 7.64 (m, 2 H) 7.97 (d,  $J=2.34$  Hz, 1 H). MS (ESI) ( $M+H$ )<sup>+</sup>: 527.

Example 27

- 5 *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-2-pyridinecarboxamide




- To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (42 mg, 0.12 mmol) in dichloromethane (5 ml) at 0°C was added diisopropylethylamine (84  $\mu$ L, 0.48 mmol) followed by 2-pyridinecarbonyl chloride, hydrochloride salt (43 mg, 0.24 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional
- 15 dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 21 mg of the TFA salt of *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-2-pyridinecarboxamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>)  $\delta$  1.32 (t,  $J=6.93$  Hz, 3 H) 1.68
- 20 (m, 6 H) 2.53 (m, 1 H) 3.42 (s, 3 H) 3.89 (m, 2 H) 3.94 (s, 2 H) 4.04 (q,  $J=6.93$  Hz, 2 H) 5.90 (s, 1 H) 6.74 (m, 2 H) 6.95 (m, 2 H) 7.09 (m, 1 H) 7.32 (m, 1 H) 7.55 (m, 1 H) 7.59 (m, 1 H) 7.76 (m, 1 H) 8.16 (s, 1 H). MS (ESI) ( $M+H$ )<sup>+</sup>: 456.

Example 28

- 25 *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide




- 56 -



To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (71 mg, 0.20 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (107  $\mu$ L, 0.61 mmol) followed by isovaleryl chloride (50  $\mu$ L, 0.41 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2 M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using reversed phase silica gel flash chromatography (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 20 mg of the TFA salt of *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>)  $\delta$  0.69 (d, *J*=6.64 Hz, 6 H) 1.25 (t, *J*=6.93 Hz, 3 H) 1.72 (m, 6 H) 1.80 (m, 1 H) 1.91 (d, *J*=6.64 Hz, 2 H) 2.64 (d, *J*=7.23 Hz, 1 H) 3.17 (s, 3 H) 3.89 (q, *J*=6.93 Hz, 2 H) 4.02 (s, 2 H) 4.11 (d, *J*=7.23 Hz, 2 H) 6.05 (s, 1 H) 6.75 (m, 2 H) 7.02 (d, *J*=8.59 Hz, 2 H) 7.62 (d, *J*=2.15 Hz, 1 H) 7.91 (d, *J*=2.15 Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 435.

**Example 29**

*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea




To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (71 mg, 0.20 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (107  $\mu$ L, 0.61 mmol) followed by isopropylisocyanate (41  $\mu$ L, 0.41

- 57 -

mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2 M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using reversed phase silica gel flash chromatography (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 18mg of the TFA salt of *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 0.93 (d, *J*=6.64 Hz, 6 H) 1.25 (t, *J*=6.96 Hz, 3 H) 1.72 (m, 4 H) 1.80 (m, 2 H) 2.66 (m, 1 H) 3.13 (s, 3 H) 3.77 (m, 1 H) 3.89 (q, *J*=6.96 Hz, 2 H) 4.02 (s, 2 H) 4.09 (d, *J*=7.23 Hz, 2 H) 6.07 (s, 1 H) 6.74 (d, *J*=8.79 Hz, 2 H) 7.01 (d, *J*=8.79 Hz, 2 H) 7.66 (d, *J*=2.34 Hz, 1 H) 7.93 (d, *J*=2.34 Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 436.

Example 30

*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,1-dimethyl-1*H*-imidazole-5-sulfonamide




To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (43 mg, 0.12 mmol) in dichloromethane (5 ml) at 0°C was added diisopropylethylamine (84 μL, 0.48 mmol) followed by the hydrochloride salt of 1-methyl 1*H*-imidazole-5-sulfonyl chloride (52 mg, 0.24 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 14 mg of the TFA salt of *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,1-dimethyl-1*H*-imidazole-5-sulfonamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 1.25 (m, 3

- 58 -

H) 1.75 (m, 6 H) 2.62 (s, 1 H) 3.25 (s, 3 H) 3.63 (m, 3 H) 3.90 (q,  $J=7.03$  Hz, 2 H) 3.99 (s, 2 H) 4.06 (d,  $J=7.23$  Hz, 2 H) 5.98 (s, 1 H) 6.75 (m, 2 H) 7.00 (d,  $J=8.59$  Hz, 2 H) 7.38 (d,  $J=1.17$  Hz, 1 H) 7.49 (d,  $J=2.34$  Hz, 1 H) 7.68 (s, 1 H) 7.84 (d,  $J=2.34$  Hz, 1 H). MS (ESI)  $(M+H)^+$  : 495

# 5 Example 31

*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,5-dimethyl-3-isoxazolecarboxamide




10

To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (57 mg, 0.16 mmol) in dichloromethane (5 ml) at 0°C was added diisopropylethylamine (84  $\mu$ L, 0.48 mmol) followed by 5-methyl-3-isoxazolecarbonyl chloride (48 mg, 0.33 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M  $\text{Na}_2\text{CO}_3$  aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with  $\text{MgSO}_4$ , filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 25 mg of the TFA salt of 3-isoxazolecarboxamide *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,5-dimethyl-3-isoxazolecarboxamide.  $^1\text{H}$  NMR (400 MHz, METHANOL- $\text{D}_4$ )  $\delta$  1.26 (m, 3 H) 1.71 (m, 6 H) 2.12 (s, 3 H) 2.61 (m, 1 H) 3.37 (s, 3 H) 3.90 (q,  $J=6.96$  Hz, 2 H) 3.99 (s, 2 H) 4.08 (d,  $J=7.23$  Hz, 2 H) 5.84 (s, 1 H) 5.99 (s, 1 H) 6.75 (m, 2 H) 7.01 (d,  $J=8.59$  Hz, 2 H) 7.63 (d,  $J=2.34$  Hz, 1 H) 7.86 (d,  $J=2.34$  Hz, 1 H). MS (ESI)  $(M+H)^+$  : 460

# 25 Example 32

*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-4-(dimethylamino)-*N*-methyl- benzamide


- 59 -



To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-amine (52 mg, 0.15 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (78 µL, 0.45 mmol) followed by 4-(dimethylamino)-benzoyl chloride (55 mg, 0.30 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 19 mg of the TFA salt of N-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]-4-(dimethylamino)-N-methylbenzamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 1.25 (t, *J*=6.93 Hz, 3 H) 1.68 (m, 6 H) 2.58 (m, 1 H) 2.76 (s, 6 H) 3.36 (s, 3 H) 3.89 (q, *J*=6.93 Hz, 2 H) 3.96 (s, 2 H) 4.03 (d, *J*=7.22 Hz, 2 H) 5.95 (s, 1 H) 6.40 (d, *J*=8.79 Hz, 2 H) 6.74 (d, *J*=8.59 Hz, 2 H) 7.00 (d, *J*=8.59 Hz, 2 H) 7.06 (d, *J*=8.79 Hz, 2 H) 7.62 (d, *J*=2.34 Hz, 1 H) 7.71 (d, *J*=2.34 Hz, 1 H). MS (ESI) (M+H)<sup>+</sup>: 498

**Example 33**

4-[[[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-yl]methylamino]sulfonyl]-benzoic acid




To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1H-pyrrolo[2,3-b]pyridin-5-amine (60 mg, 0.17 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (118 µL, 0.68 mmol) followed by 4-(chlorosulfonyl)-benzoic acid (76

- 60 -

mg, 0.34 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 12 mg of the TFA salt of 4-[[[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]methylamino]sulfonyl]-benzoic acid. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 1.25 (t, *J*=7.03 Hz, 3 H) 1.72 (m, 6 H) 2.60 (m, 1 H) 3.41 (s, 3 H) 3.90 (q, *J*=7.03 Hz, 2 H) 4.03 (s, 2 H) 4.10 (m, 2 H) 5.98 (s, 1 H) 6.77 (d, *J*=8.59 Hz, 2 H) 7.03 (d, *J*=8.59 Hz, 2 H) 7.28 (m, 2 H) 7.54 (m, 2 H) 8.08 (s, 1 H) 8.19 (s, 1 H). MS (ESI) (M+H)<sup>+</sup>: 535

Example 34

*N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-2-nitro- benzenesulfonamide



15

To a solution of 1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-amine (52 mg, 0.15 mmol) in dichloromethane (5 mL) at 0°C was added diisopropylethylamine (78 μL, 0.45 mmol) followed by 2-nitro- benzenesulfonyl chloride (65 mg, 0.30 mmol). The mixture was stirred overnight at room temperature. The reaction was then quenched by 2M Na<sub>2</sub>CO<sub>3</sub> aqueous solution (10 mL). The phases were separated and the aqueous phase was back-extracted with additional dichloromethane (10 mL). The organic phases were combined, dried with MgSO<sub>4</sub>, filtered and concentrated *in vacuo*. The residue was purified using HPLC (10 to 70% [0.1% TFA in AcCN solution] in 0.1% TFA aqueous solution) to provide 21 mg of the TFA salt of *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-2-nitro- benzenesulfonamide. <sup>1</sup>H NMR (400 MHz, METHANOL-D<sub>4</sub>) δ 1.25 (t, *J*=6.93 Hz, 3 H) 1.72 (m, 6 H) 2.61 (m, 1 H) 3.28 (s, 3 H) 3.88 (q, *J*=6.93 Hz, 2 H) 3.98 (s, 2 H) 4.07 (d, *J*=7.23 Hz,

25


- 61 -

2 H) 5.97 (s, 1 H) 6.73 (d,  $J=8.79$  Hz, 2 H) 6.99 (d,  $J=8.79$  Hz, 2 H) 7.47 (m, 2 H) 7.53 (d,  $J=2.34$  Hz, 1 H) 7.65 (m, 2 H) 7.82 (d,  $J=2.34$  Hz, 1 H). MS (ESI)  $(M+H)^+$ : 536

- 62 -

What is claimed is:

1. A compound of formula I or a pharmaceutically acceptable salt thereof:



**I**

5 wherein

$R^1$  is a  $C_{1-12}$  group;

$X$  is a  $C_{1-10}$  divalent group that separates groups connected thereto by one or two saturated carbons;

$Ar$  is  $C_{4-12}$  divalent aromatic group;

10  $R^2$  is optionally substituted  $C_{1-6}$ hydrocarbaryl, optionally substituted  $C_{6-10}$ aryl, or optionally substituted  $C_{3-6}$ heteroaryl;

$R^3$  is a  $C_{1-12}$  group, wherein the atom of  $R^3$  that is directly connected to the six-membered ring of formula I is a nitrogen, or an unsaturated carbon, wherein the unsaturated carbon is connected to an oxygen through a double bond; and

15  $R^a$  and  $R^b$  are  $-R$ ,  $-NO_2$ ,  $-OR$ ,  $-Cl$ ,  $-Br$ ,  $-I$ ,  $-F$ ,  $-CF_3$ ,  $-C(=O)R$ ,  $-C(=O)OH$ ,  $-NH_2$ ,  $-SH$ ,  $-NHR$ ,  $-NR_2$ ,  $-SR$ ,  $-SO_3H$ ,  $-SO_2R$ ,  $-S(=O)R$ ,  $-CN$ ,  $-OH$ ,  $-C(=O)OR$ , or  $-NRC(=O)R$ , wherein  $R$  is independently  $-H$  or  $C_{1-6}$  hydrocarbaryl.

2. A compound as claimed in claim 1, wherein


20  $R^1$  is optionally substituted  $C_{1-10}$  hydrocarbaryl; optionally substituted  $C_{1-10}$ acyl; optionally substituted  $C_{4-8}$ heteroaryl- $C(=O)-$ ;  $R^4R^5N-C_{1-6}$ alkyl;  $R^4R^5NC(=O)-C_{1-6}$ alkyl;  $R^4O-C_{1-6}$ alkyl;  $R^4OC(=O)-C_{1-6}$ alkyl;  $R^4C(=O)-C_{1-6}$ alkyl;  $R^4C(=O)NR^4-C_{1-6}$ alkyl;  $R^4R^5NSO_2-C_{1-6}$ alkyl;  $R^4CSO_2N(R^5)-C_{1-6}$ alkyl;  $R^4R^5NC(=O)N(R^6)-C_{1-6}$ alkyl;  $R^4R^5NSO_2N(R^6)-C_{1-6}$ alkyl; optionally substituted aryl- $C_{1-6}$ alkyl; optionally substituted aryl- $C(=O)-C_{1-6}$ alkyl; optionally substituted heterocyclyl- $C_{1-6}$ alkyl; optionally substituted heterocyclyl- $C(=O)-C_{1-6}$ alkyl; and  $C_{1-10}$ hydrocarbaryl-amino;

25

- 63 -

wherein  $R^4$ ,  $R^5$  and  $R^6$  are independently selected from -H,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl, or a divalent  $C_{1-6}$ group that together with another divalent  $C_{1-6}$ group forms a portion of a ring;

$R^3$  is selected from:



5

wherein

$R^7$  is selected from -H, optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl;

10  $R^{10}$ ,  $R^{11}$ ,  $R^{12}$  and  $R^{13}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl; and  $R^a$  and  $R^b$  are hydrogen.

15 3. A compound as claimed claim 1,

wherein  $R^1$  is selected from  $C_{1-8}$ alkyl;  $C_{2-8}$ alkenyl;  $C_{2-8}$ alkynyl; optionally substituted aryl- $C_{1-6}$ alkyl;  $R^4R^5NC_{1-6}$ alkyl;  $R^4OC_{1-6}$ alkyl;  $C_{3-6}$ cycloalkyl- $C_{1-6}$ alkyl; optionally substituted  $C_{3-6}$ heterocycloalkyl- $C_{1-6}$ alkyl;  $C_{1-6}$ alkyl- $C_{6-8}$ aryl;  $C_{1-6}$ alkyl-C(=O)-;  $C_{6-8}$ aryl-C(=O)-;  $C_{3-8}$ heteroaryl-C(=O)-; or optionally substituted  $C_{3-6}$ heteroaryl- $C_{1-6}$ alkyl;

20 wherein  $R^2$  is selected from  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyl substituted by at least one fluorine,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkenyl substituted by at least one fluorine,  $C_{2-6}$ alkynyl,  $C_{2-6}$ alkynyl substituted by at least one fluorine, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$ aryl, and optionally substituted  $C_{3-6}$ heteroaryl;


25  $R^4$ ,  $R^5$  and  $R^6$  are independently selected from the group consisting of -H,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl, and a divalent  $C_{1-6}$ group that together with another divalent  $C_{1-6}$ group forms a portion of a ring;



- 64 -

X is selected from the group consisting of  $-\text{NR}^6-$ ,  $-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}=\text{CH}-$ ,  $-\text{O}-$ ,  $-\text{C}(\text{R}^8)(\text{R}^9)-$ , and  $-\text{S}(\text{O})_q-$ , wherein q is 0, 1 or 2, wherein  $\text{R}^8$  and  $\text{R}^9$  are independently  $\text{C}_{1-6}$ alkyl,  $\text{C}_{2-6}$ alkenyl,  $\text{C}_{2-6}$ alkynyl,  $\text{C}_{1-6}$ alkoxy,  $-\text{OH}$ , or  $-\text{H}$ ; at most one of  $\text{R}_8$  and  $\text{R}_9$  is  $-\text{OH}$ ;

$\text{R}^3$  is selected from:



5

wherein

$\text{R}^7$  is selected from  $-\text{H}$ , optionally substituted  $\text{C}_{1-6}$ alkyl, optionally substituted  $\text{C}_{2-6}$ alkenyl, optionally substituted  $\text{C}_{2-6}$ alkynyl, optionally substituted  $\text{C}_{3-6}$ cycloalkyl, optionally substituted  $\text{C}_{6-10}$  aryl, or optionally substituted  $\text{C}_{3-6}$ heteroaryl;

- 10  $\text{R}^{10}$ ,  $\text{R}^{11}$ ,  $\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from optionally substituted  $\text{C}_{1-6}$ alkyl, optionally substituted  $\text{C}_{2-6}$ alkenyl, optionally substituted  $\text{C}_{2-6}$ alkynyl, optionally substituted  $\text{C}_{3-6}$ cycloalkyl, optionally substituted  $\text{C}_{6-10}$  aryl, or optionally substituted  $\text{C}_{3-6}$ heteroaryl; and  $\text{R}^a$  and  $\text{R}^b$  are hydrogen.

- 15 4. A compound as claimed in claim 3, wherein


$\text{R}^1$  is selected from  $\text{C}_{1-6}$ alkyl;  $\text{C}_{2-6}$ alkenyl;  $\text{C}_{2-6}$ alkynyl; optionally substituted  $\text{C}_{3-6}$ cycloalkylmethyl; optionally substituted  $\text{C}_{3-6}$ heterocycloalkylmethyl;

X is  $-\text{CH}_2-$ ;

Ar is phenylene or pyridylene;

- 20  $\text{R}^2$  is selected from  $-\text{CH}_3$ ,  $-\text{CH}_2\text{CH}_3$ ,  $-\text{CH}(\text{CH}_3)_2$ ,  $-\text{CH}_2\text{CF}_3$ ,  $\text{CF}_3$ , cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyridyl and phenyl; and

$\text{R}^3$  is selected from



- 25 wherein,  $\text{R}^7$  is selected from  $-\text{H}$  and methyl;  $\text{R}^{10}$  and  $\text{R}^{11}$  are independently selected from optionally substituted  $\text{C}_{1-6}$ alkyl, optionally substituted  $\text{C}_{2-6}$ alkenyl, optionally substituted

- 65 -

C<sub>2-6</sub>alkynyl, optionally substituted C<sub>3-6</sub>cycloalkyl, optionally substituted C<sub>6-10</sub> aryl, or optionally substituted C<sub>3-6</sub>heteroaryl.

5. A compound as claimed in claim 3, wherein


5 R<sup>1</sup> is selected from C<sub>1-6</sub>alkyl; C<sub>2-6</sub>alkenyl; C<sub>2-6</sub>alkynyl; optionally substituted C<sub>3-6</sub>cycloalkylmethyl; optionally substituted C<sub>3-6</sub>heterocycloalkylmethyl;

X is -CH<sub>2</sub>-;

Ar is selected from the group consisting of an optionally substituted *para*-arylene; an optionally substituted a six-membered *para*-heteroarylene;

10 R<sup>2</sup> is selected from -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CF<sub>3</sub>, CF<sub>3</sub>, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyridyl and phenyl; and

R<sup>3</sup> is selected from:



15 wherein, R<sup>7</sup> is selected from -H and methyl; R<sup>10</sup> and R<sup>11</sup> are selected from optionally substituted C<sub>1-6</sub>alkyl, optionally substituted C<sub>2-6</sub>alkenyl, optionally substituted C<sub>2-6</sub>alkynyl, optionally substituted C<sub>3-6</sub>cycloalkyl, optionally substituted C<sub>6-10</sub> aryl, or optionally substituted C<sub>3-6</sub>heteroaryl.

6. A compound as claimed in claim 3, wherein


20 R<sup>1</sup> is selected from optionally substituted C<sub>3-6</sub>cycloalkylmethyl; and optionally substituted C<sub>3-6</sub>heterocycloalkylmethyl;

X is -CH<sub>2</sub>-;

Ar is *para*-phenylene or *para*-pyridylene;

R<sup>2</sup> is methyl, or ethyl; and

25 R<sup>3</sup> is selected from



wherein, R<sup>7</sup> is selected from -H and methyl; R<sup>10</sup> and R<sup>11</sup> are selected from C<sub>1-6</sub>alkyl, C<sub>3-6</sub>cycloalkyl, phenyl optionally substituted with halogen, nitro, C<sub>1-3</sub>alkyl, -COOR<sup>14</sup>, -OH,

cyano, trifluormethyl, C<sub>1-3</sub>alkyloxy; C<sub>3-6</sub>heteroaryl optionally substituted with halogen, nitro, C<sub>1-3</sub>alkyl, -COOR<sup>14</sup>, -OH, cyano, trifluormethyl, C<sub>1-3</sub>alkyloxy, wherein R<sup>14</sup> is a C<sub>1-3</sub>alkyl.

7. A compound selected from:

- 5 1) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide;
- 2) *N*-[1-(cyclohexylmethyl)-2-[(3-methoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide;
- 10 3) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N*-(1-methylethyl)-urea;
- 4) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide;
- 5) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2-dimethyl-propanamide;
- 15 6) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-cyclopropanecarboxamide;
- 7) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2,2-trimethyl-propanamide;
- 8) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,*N*'-diethyl-*N*-methyl-urea;
- 20 9) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,5-dimethyl-3-isoxazolecarboxamide;
- 10) *N*-[1-(cyclohexylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2-fluoro-*N*-methyl-benzamide;
- 25 11) *N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide;
- 12) [1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-, 1-methylethyl ester carbamic acid;
- 13) *N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2,2-trimethyl-propanamide;
- 30 14) *N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide;

- 15) *N*-[1-(cyclohexylmethyl)-2-[(5-ethoxy-2-pyridinyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea;
- 16) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide;
- 5 17) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,6-difluoro-*N*-methyl-benzenesulfonamide;
- 18) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-cyclobutanecarboxamide;
- 19) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,5-difluoro-*N*-methyl-benzamide;
- 10 20) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2-dimethyl-propanamide;
- 21) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,2,2-trimethyl-propanamide;
- 15 22) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea;
- 23) *N*-[2-[(4-ethoxyphenyl)methyl]-1-[(tetrahydro-2*H*-pyran-4-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide;
- 24) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,2-dimethyl-propanamide;
- 20 25) [1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-, methyl ester carbamic acid;
- 26) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-2,6-difluoro-*N*-methyl-benzenesulfonamide;
- 25 27) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-2-pyridinecarboxamide;
- 28) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,3-dimethyl-butanamide;
- 29) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-*N'*-(1-methylethyl)-urea;
- 30 30) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,1-dimethyl-1*H*-imidazole-5-sulfonamide;

- 68 -

- 31) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-4-(dimethylamino)-*N*-methyl- benzamide;
- 32) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*,5-dimethyl-3-isoxazolecarboxamide;
- 5 33) 4-[[[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]methylamino]sulfonyl]-benzoic acid;
- 34) *N*-[1-(cyclobutylmethyl)-2-[(4-ethoxyphenyl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl]-*N*-methyl-2-nitro-benzenesulfonamide; and pharmaceutically acceptable salts thereof.

10

8. A compound according to any one of claims 1-7 for use as a medicament.

9. The use of a compound according to any one of claims 1-7 in the manufacture of a medicament for the therapy of pain.

15

10. The use of a compound according to any one of claims 1-7 in the manufacture of a medicament for the treatment of immune cancer.

20

11. The use of a compound according to any one of claims 1-7 in the manufacture of a medicament for the treatment of multiple sclerosis, Parkinson's disease, Huntington's chorea, Alzheimer's disease, anxiety disorders, gastrointestinal disorders or cardiovascular disorders.

12. A pharmaceutical composition comprising a compound according to any one of claims 1-7 and a pharmaceutically acceptable carrier.


25

13. A method for the therapy of pain in a warm-blooded animal, comprising the step of administering to said animal in need of such therapy a therapeutically effective amount of a compound according to any one of claims 1-7.

30


14. A method for preparing a compound of formula II,

- 69 -




comprising the steps of

a) reacting a compound of formula III,



5

with a base having a pKa more than 20;



b) reacting a product formed in step a) with a compound of formula IV,  
to form the compound of formula II,

wherein


- 10  $R^1$  is optionally substituted  $C_{1-10}$  hydrocarbyl; optionally substituted  $C_{1-10}$  acyl;  
optionally substituted  $C_{4-8}$  heteroaryl- $C(=O)-$ ;  $R^4R^5N-C_{1-6}$  alkyl;  $R^4R^5NC(=O)-C_{1-6}$  alkyl;  $R^4O-$   
15  $C_{1-6}$  alkyl;  $R^4OC(=O)-C_{1-6}$  alkyl;  $R^4C(=O)-C_{1-6}$  alkyl;  $R^4C(=O)NR^4-C_{1-6}$  alkyl;  $R^4R^5NSO_2-C_{1-6}$   
alkyl;  $R^4CSO_2N(R^5)-C_{1-6}$  alkyl;  $R^4R^5NC(=O)N(R^6)-C_{1-6}$  alkyl;  $R^4R^5NSO_2N(R^6)-C_{1-6}$  alkyl;  
optionally substituted aryl- $C_{1-6}$  alkyl; optionally substituted aryl- $C(=O)-C_{1-6}$  alkyl; optionally  
20 substituted heterocyclyl- $C_{1-6}$  alkyl; optionally substituted heterocyclyl- $C(=O)-C_{1-6}$  alkyl; and  
 $C_{1-10}$  hydrocarbylamino;

wherein  $R^4$ ,  $R^5$  and  $R^6$  are independently selected from -H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, or a divalent  $C_{1-6}$  group that together with another divalent  $C_{1-6}$  group forms a portion  
20 of a ring;

$R^2$  is optionally substituted  $C_{1-6}$  hydrocarbyl, optionally substituted  $C_{6-10}$  aryl, or  
optionally substituted  $C_{3-6}$  heteroaryl;

$R^3$  is selected from:

- 70 -



wherein

- $R^7$  is selected from -H, optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl;

$R^{10}$ ,  $R^{11}$ ,  $R^{12}$  and  $R^{13}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl; and  $R^c$  is  $C_{1-4}$ alkyl.

10


15. A process as claimed in claim 14, wherein the base is t-butyl lithium;

$R^1$  is selected from  $C_{1-6}$ alkyl;  $C_{2-6}$ alkenyl;  $C_{2-6}$ alkynyl; optionally substituted  $C_{3-6}$ cycloalkylmethyl; optionally substituted  $C_{3-6}$ heterocycloalkylmethyl;

15

$R^2$  is selected from -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CF<sub>3</sub>, CF<sub>3</sub>, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyridyl and phenyl; and

$R^3$  is selected from:




- wherein,  $R^7$  is selected from -H and methyl;  $R^{10}$  and  $R^{11}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl.


20

16. A process for preparing a compound of formula V,


- 71 -

V

comprising the step of reacting a compound of formula VI,

VI

with a compound of formula VII,

VII

5

to form the compound of formula V,

wherein

R¹ is optionally substituted C<sub>1-10</sub> hydrocarbyl; optionally substituted C<sub>1-10</sub>acyl; optionally substituted C<sub>4-8</sub>heteroaryl-C(=O)-; R⁴R⁵N-C<sub>1-6</sub>alkyl; R⁴R⁵NC(=O)-C<sub>1-6</sub>alkyl; R⁴O-C<sub>1-6</sub>alkyl; R⁴OC(=O)-C<sub>1-6</sub>alkyl; R⁴C(=O)-C<sub>1-6</sub>alkyl; R⁴C(=O)NR⁴-C<sub>1-6</sub>alkyl; R⁴R⁵NSO<sub>2</sub>-C<sub>1-6</sub>alkyl; R⁴CSO<sub>2</sub>N(R⁵)-C<sub>1-6</sub>alkyl; R⁴R⁵NC(=O)N(R⁶)-C<sub>1-6</sub>alkyl; R⁴R⁵NSO<sub>2</sub>N(R⁶)-C<sub>1-6</sub>alkyl; optionally substituted aryl-C<sub>1-6</sub>alkyl; optionally substituted aryl-C(=O)-C<sub>1-6</sub>alkyl; optionally substituted heterocyclyl-C<sub>1-6</sub>alkyl; optionally substituted heterocyclyl-C(=O)-C<sub>1-6</sub>alkyl; and C<sub>1-10</sub>hydrocarbylamino;


15 wherein R⁴, R⁵ and R⁶ are independently selected from -H, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, or a divalent C<sub>1-6</sub>group that together with another divalent C<sub>1-6</sub>group forms a portion of a ring;

R² is optionally substituted C<sub>1-6</sub>hydrocarbyl, optionally substituted C<sub>6-10</sub>aryl, or optionally substituted C<sub>3-6</sub>heteroaryl;

20 R³ is selected from:



- 72 -



wherein

$R^7$  is selected from -H, optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl;

$R^{10}$ ,  $R^{11}$ ,  $R^{12}$  and  $R^{13}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl;

Y is CH or N; and


$R^c$  is  $C_{1-4}$ alkyl.

17. A process as claimed in claim 16, wherein

$R^1$  is selected from  $C_{1-6}$ alkyl;  $C_{2-6}$ alkenyl;  $C_{2-6}$ alkynyl; optionally substituted  $C_{3-6}$ cycloalkylmethyl; optionally substituted  $C_{3-6}$ heterocycloalkylmethyl;

$R^2$  is selected from -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CF<sub>3</sub>, CF<sub>3</sub>, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyridyl and phenyl; and

$R^3$  is selected from:



wherein,  $R^7$  is selected from -H and methyl;  $R^{10}$  and  $R^{11}$  are independently selected from optionally substituted  $C_{1-6}$ alkyl, optionally substituted  $C_{2-6}$ alkenyl, optionally substituted  $C_{2-6}$ alkynyl, optionally substituted  $C_{3-6}$ cycloalkyl, optionally substituted  $C_{6-10}$  aryl, or optionally substituted  $C_{3-6}$ heteroaryl.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2004/000472

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 471/04, A61K 31/437, A61P 25/04  
According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CHEM.ABS.DATA, WPI DATA

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 0158869 A2 (BRISTOL-MYERS SQUIBB COMPANY), 16 August 2001 (16.08.2001)  --	1-17
A	WO 9822457 A1 (AMGEN INC.), 28 May 1998 (28.05.1998)  -- -----	1-17

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

5 July 2004

Date of mailing of the international search report

08-07-2004

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

CAROLINA GÓMEZ LAGERLÖF/BS  
Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

Information on patent family members

30/04/2004

International application No.

PCT/SE 2004/000472

WO	0158869	A2	16/08/2001	AU	3495801	A	20/08/2001
				CA	2399791	A	16/08/2001
				EP	1254115	A	06/11/2002
				JP	2004502642	T	29/01/2004
				US	6653304	B	25/11/2003
				US	2002119972	A	29/08/2002
<hr/>							
WO	9822457	A1	28/05/1998	AT	264318	T	15/04/2004
				AU	734841	B	21/06/2001
				AU	5265998	A	10/06/1998
				CA	2271767	A	28/05/1998
				CN	1246856	A	08/03/2000
				EP	0948495	A,B	13/10/1999
				HU	9903330	A	28/03/2000
				IL	129928	D	00/00/0000
				JP	2001506980	T	29/05/2001
				KR	2000057137	A	15/09/2000
				US	6180643	B	30/01/2001
				US	6440973	B	27/08/2002
				US	6605634	B	12/08/2003
				US	2003096819	A	22/05/2003

---

**INTERNATIONAL SEARCH REPORT****International application No.**  
**PCT/SE 2004/000472****Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **13**  
because they relate to subject matter not required to be searched by this Authority, namely:  
**see extra sheet**
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**

International application No.  
**PCT/SE 2004/000472**

**Box II.1**

Claim 13 relates to methods of treatment of the human or animal body by surgery or by therapy or diagnostic methods practiced on the human or animal body (Rule 39.1(iv)). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compounds or compositions